

REFERENCES

- AGTERBERG, F.P. & GRADSTEIN, F.M. (1999). – The RASC method for ranking and scaling of biostratigraphic events. – *Earth Science Reviews*, 46 (1-4), 1-27.
- AKIMOTO, K., HATTORI, M., UEMATSU, K. & KATO, C. (2001). – The deepest living foraminifera, Challenger Deep, Mariana Trench. – *Marine Micropaleontology*, 42, 95-97.
- AKSU, A.E., ABRAJANO, T., MUDIE, P.J. & JAŞAR, D. (1999). – Organic geochemical and palynological evidence for terrigenous origin of the organic matter in Aegean Sea sapropel S1. – *Marine Geology*, 153, 308-318.
- ALTENBACH, A., PFLAUMANN, U., SCHIEBEL, R., THIES, A., TIMM, S. & TRAUTH, M. (1999). – Scaling percentages and distributional patterns of benthic foraminifera with flux rates of organic carbon. – *Journal of Foraminiferal Research*, 29 (3), 173-185.
- ALTENBACH, A.V., UNSOLD, G. & WALGER, E. (1988) – The hydrodynamic Environment of *Saccorhiza ramosa* (Brady). – *Meyniana*, 40, 119-132.
- ALTH, A. (1850). – Geognostische-paläntologische Beschreibung der nächsten Umgebung von Lemberg. – *Abhandlungen Naturwissenschaften Wien*, 3, 171-284.
- APPLIN, E.R., LOEBLICH, A.R. & TAPPAN, H. (1950). – Two new Lower Cretaceous luluolid foraminifera. – *Journal of the Washington Academy of Sciences, Washington, D.C.*, 40, 75-79.
- ARAPOVA, N.D. & SULEYMANOV, I.S. (1966). – O foraminiferakh iz Konyakskikh otlozhenii zapadnogo Uzbekistan i Kyzylkumov [On the foraminifera of the Coniacian deposits of western Uzbekistan and the Kyzyl-Kum]. – Tashkentskii Gosudarstvennyy Universitet im. V.I. Lenina, 273, 121-127.
- ARTHUR, M.A. (1979). – *Sedimentologic and geochemical studies of Late Cretaceous and Paleocene pelagic sedimentary rocks, Part 1: The Gubbio sequence*. Ph.D. Dissertation, Princeton University, 1-171.
- ARTHUR, M.A. & FISCHER, A.G. (1977). – Upper Cretaceous – Paleocene magnetic stratigraphy at Gubbio, Italy: 1 – Lithostratigraphy and sedimentology. – *Geological Society of America Bulletin*, 88 (3), 367-371.
- ARTHUR, M.A. & PREMOLI-SILVA, I. (1982). – Development of widespread organic carbon-rich strata in Mediterranean Tethys. – In: Schlanger, S.O. & Cita, M.B. (eds), *Nature and Origin of Cretaceous carbon-rich facies*. Academic Press, New York, 9-54.
- AUBERT, J. & BERGGREN, W.A. (1976). – Paleocene benthic foraminiferal biostratigraphy and paleoecology of Tunisia. – *Bulletin des Centre Rech. Pau – SNPA*, 10, 379-469.
- AVNEMELICH, M. (1952). – A revision of the tubular Monothalamia. – *Contributions from the Cushman Foundation for Foraminiferal Research*, 3, 60-68.
- BAK, K. (2000). – Biostratigraphy of deep-water agglutinated Foraminifera in Scaglia Rossa-type deposits of the Pieniny Klippen Belt, Carpathians, Poland. – In: Hart, M.B., Kaminski, M.A. & Smart, C.W. (eds), *Proceedings of the Fifth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 7, 15-40.
- BAK, K. (2004). – Deep-water agglutinated foraminiferal changes across the Cretaceous/Tertiary and Paleocene/Eocene transitions in the deep flysch environment; eastern Outer Carpathians, (Bieszczady Mts., Poland). – In: Bubík, M. & Kaminski, M.A., (eds), *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 8, 1-56.
- BALKWELL, F.P. & WRIGHT, J. (1885). – Report on some Recent Foraminifera found off the coast of Dublin and the Irish Sea. – *Transactions of the Royal Irish Academy*, 28, 317-372.
- BANDY, O.L. (1963). – Larger living foraminifera of the continental borderland off southern California. – *Contributions from the Cushman Foundation for Foraminiferal Research*, 14, 121-126.
- BANDY, O.L. & ECHOLS, R.J. (1964). – Antarctic foraminiferal zonation. – In: Lee, M.O. (Ed.), *Biology of the Antarctic Seas*. Antarctic Research Series, 1, 73-91.
- BANDY, O.L. & RODOLFO, K.S. (1964). – Distribution of foraminifera and sediments, Peru-Chile Trench area. – *Deep Sea Research*, 11, 817-837.
- BANNER, F.T. (1966). – Morfologiya, klassifikatsiya i stratigraficheskoye znachenie spirotsiklinid. – *Voprosy Mikropaleontologii*, 10, 201-224.
- BANNER, F.T. (1970). – A synopsis of the Spirocyclinidae. – *Revista Española de Micropaleontología*, 2, 243-290.
- BANNER, F.T. & PEREIRA, C.P.G. (1981). – Some biserial and triserial agglutinated smaller foraminifera: their wall structure and its significance. – *Journal of Foraminiferal Research*, 11, 85-117.

- BANNER, F.T., DESAI, D. (1985). – The genus *Clavulinoides* Cushman emended and the new Cretaceous genus *Clavulinopsis*. – *Journal of Foraminiferal Research*, 15, 79-90.
- BARKER, R.W. (1960). – *Taxonomic notes on the species figured by H.R. Brady in his report on the foraminifera dredged by H.M.S. Challenger during the years 1873–1876*. – Society of Economic Paleontologists and Mineralogists Special Publication, 9, 238 pp. Tulsa, Oklahoma.
- BARTENSTEIN, H. (1952). – Taxonomische Bemerkungen zu den *Ammobaculites*, *Haplophragmium*, *Lituola* und verwandten Gattungen (For.). – *Senckenbergiana*, 33, 313-342.
- BARTENSTEIN, H. (1954). – Revision von Berthelin's Memoire 1880 über Alb-Foraminiferen von Montcley. – *Senckenbergiana Lethaea*, 35, 37-50.
- BARTENSTEIN, H. & BRAND, E. (1951). – Mikropaläontologie Untersuchungen zur Stratigraphie des nordwestdeutschen Valendis. – *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, 485, 239-336.
- BATJES, D.A.J. (1958). – Foraminifera of the Oligocene of Belgium. – *Institut Royal des Sciences Naturelles de Belgique, Memoire*, 143, 1-188.
- BALDAUF, J.G., CLEMENT, B., AKSU, A.E., DEVERNAL, A., FIRTH, J.V., HALL, F., HEAD, M.J., JARRARD, R., KAMINSKI, M.A., LAZARUS, D.B., MONJANEL, A.-L., BERGGREN, W.A., GRADSTEIN, F.M., KNUTTTEL, S., MUDIE, P. & RUSSEL, M. (1989). – Magnetostratigraphic and biostratigraphic synthesis of Ocean Drilling Program Leg 105; Labrador Sea and Baffin Bay. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 105, 935-956.
- BECK, R.S. (1943). – Eocene foraminifera from Cowlitz River, Lewis County, Washington. – *Journal of Paleontology*, 17 (6), 584-614.
- BECKMANN, J.-P. (1960). – Distribution of benthonic foraminifera at the Cretaceous Tertiary boundary of Trinidad (West Indies). – International Geological Congress, 21 Session, Norden, 5, 57-69.
- BECKMANN, J.-P. (1978). – Late Cretaceous smaller benthic foraminifera from Sites 363 and 364, DSDP Leg 40, Southeast Atlantic Ocean. – *Initial Reports of the Deep Sea Drilling Project*, 40, 759-781.
- BECKMANN, J.-P. (1991). – New taxa of foraminifera from the Cretaceous and basal Tertiary of Trinidad, West Indies. – *Eclogae Geologicae Helveticae*, 84 (3), 819-835.
- BECKMANN, J.-P. (1994). – Late Albian to Early Eocene. – In: Bolli, H.M., Beckmann, J.P. & Saunders, J.B. (eds), *Benthic foraminiferal biostratigraphy of the south Caribbean region*. Cambridge University Press, pp. 47-227.
- BECKMANN, J.-P., BOLLI, H.M., KLEBOTH, P. & PROTO-DECIMA, F. (1982). – Micropaleontology and biostratigraphy of the Campanian to Paleocene of the Monte Giglio, Bergamo Province, Italy. – *Memorie di Scienze Geologiche*, 35, 91-172.
- BENDER, H. (1989). – Gehäuseaufbau, Gehäusegenese, und Biologie agglutiniertes Foraminiferen (Sarcodina, Textulariina). – *Jahrbuch der Geologischen Bundesanstalt, Wien*, 133, 259-347.
- BENDER, H. (1995). – Test structure and classification in agglutinated foraminifera. – In: Kaminski, M.A., Geroch, S. & Gasinski, M.A. (eds), *Proceedings of the Fourth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 3, 27-70.
- BERGER, W.H. & WINTERER, E.L. (1974). – Plate stratigraphy and the fluctuating carbonate line. In: Hsü, K.J. & Jenkyns, H.C. (eds), *Pelagic sediments: On land and under the sea*. International Association of Sedimentologists Special Publication, 1, 11-98. Blackwell, Oxford.
- BERGGREN, W.A. (1972). – Cenozoic biostratigraphy and paleobiogeography of the North Atlantic. – *Initial Reports of the Deep Sea Drilling Project*, 12, 965-1001.
- BERGGREN, W.A. & KAMINSKI, M.A. (1990). – Abyssal Agglutinates: Back to Basics. – In: Hemleben, C., Kaminski, M.A., Kuhnt, W. & Scott, D.B. (eds), *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of Agglutinated Foraminifera*, NATO ASI Series C327, Kluwer Acad. Publ. 53-76.
- BERGGREN, W.A. & OLSSON, R.K. (1986). – North Atlantic Mesozoic and Cenozoic paleoceanography. – In: Vogt, P.R. & Tucholke, B.E. (eds), *The Geology of North America, Volume M, The North Atlantic Region*. Geological Society of America, 565-587.
- BERGGREN, W.A. & SCHNITKER, D. (1983). – Cenozoic marine environments in the North Atlantic and Norwegian-Greenland Sea. In: BOTT, M.H.P., SAXOV, S., TALWANI, M., THEIDE, J. (eds), *Structure and development of the Greenland- Scotland Ridge - New methods and concepts*. NATO Conference Series IV, v. 8, 495-548.
- BERMÚDEZ, P. & DE RIVERO, F.C. (1963). – *Estudio sistemático de los Foraminíferos Quitinosos, Mikrogranulares y Arenaceos*. – Universidad Central de Venezuela, Ediciones de la Biblioteca, 398 pp.
- BERMÚDEZ, P. & FUENMAYOR, A.N. (1966). – Consideraciones sobre los sedimentos del Mioceno Medio al Reciente de las Costas Central y oriental de Venezuela. Segunda parte. Los foraminíferos bentónicos. – *Boletín de Geología*, 7 (14), 413-611 + 4 pls.

- BERMÚDEZ, P. & GAMEZ, H.A. (1966). – Estudio paleontológico de una sección del Eoceno. – *Memoria de la Sociedad de Ciencias Naturales La Salle-Tomo*, 26, 205-259.
- BERTHELIN, G. (1880). – Mémoire sur les foraminifères fossiles de l'étage Albien de Montcley (Doubs). – *Mémoires de la Société Géologique de France, Series*, 3, 1, 1-84.
- BERRY, E.W. (1928). – The smaller foraminifera of the middle Lobitos shales of northwestern Peru. – *Eclogae Geologicae Helvetiae*, 21, 390-405.
- BOERSMA, A. (1985). – Biostratigraphy and biogeography of Tertiary bathyal benthic foraminifers: Tasman Sea, Coral Sea, and on the Chatam Rise (Deep Sea Drilling Project, Leg 90). – *Initial Reports of the Deep Sea Drilling Project*, 90, 961-1035.
- BOGDANOVICH, A.K. (1963). – Novye predstaviteli foraminifer s khaltsedonovoy stenкой iz tretichnykh otlozhenii severnogo Kavkaza i Kryma [New representatives of foraminifera with a chalcedony wall from the Tertiary deposits of the northern Caucasus and Crimea]. – *Voprosy Mikropaleontologii*, 7, 150-157.
- BOLLI, H.M. (1957). – Planktonic foraminifera from the Eocene Navet and San Fernando formations of Trinidad, B.W.I. – *US National Museum Bulletin*, 215, 155-172.
- BOLLI, H.M. (1994). – The Falcon Basin. – In: Bolli, H.M., Beckmann, J.P. & Saunders, J.B. (eds), *Benthic foraminiferal biostratigraphy of the south Caribbean region*. Cambridge University Press, pp. 277-303.
- BORNMALM, L., CORLISS, B.H. & TEDESCO, K. (1997). – Laboratory observations of rates and patterns of movement of continental margin benthic foraminifera. – *Marine Micropaleontology*, 29 (3-4), 175-184.
- BRADY, H.B. (1878). – On the reticularian and radiolarian rhizopoda (Foraminifera and Polycystina) of the North-Polar Expedition of 1875, 1876. – *Annals and Magazine of Natural History, series* 5, 1, 425.
- BRADY, H.B. (1879). – Notes on some of the reticularian Rhizopoda of the "CHALLENGER" Expedition; Part I. On new or little known Arenaceous types. – *Quarterly Journal of Microscopical Sciences*, 19, 20-67 + 3 pl.
- BRADY, H.B. (1881a). – Notes on some of the reticularian Rhizopoda on the "CHALLENGER" Expedition; Part III. – *Quarterly Journal of Microscopical Sciences*, 21, 31-71.
- BRADY, H.B. (1881b). – Über einige arktische Tiefsee-Foraminiferen gesammelt während der Österreichisch-ungarischen Nordpol-Expedition in den Jahren 1872-74. – *Denkschriften der D. K. Akademie Wissenschaften Wien*, 43, 91-110.
- BRADY, H.B. (1884). – Report on the foraminifera dredged by H.M.S. CHALLENGER during the years 1873-1876. – In: MURRAY, J. (ed.), *Reports of the scientific results of the voyage of the H.M.S. Challenger. Zoology*, 9, 1-814.
- BRAGA, G., DI BIASE, R., GRÜNIG, A. & PROTO DECIMA, F. (1975). – Foraminiferi bentonici del Paleocene ed Eocene della Sezione di Possagno. – *Schweizerische Paläontologische Abhandlungen*, 97, 85-111 + 6 pls.
- BRALOWER, T.J. (2002). – Evidence for surface-water oligotrophy during the late Paleocene Thermal Maximum: Nannofossil assemblage data from Ocean Drilling Program Site 690, Maud Rise, Weddell Sea. – *Paleoceanography*, 17, paper no. 10.1029/20001PA00062.
- BRATU, E. (1972). – Excursion J. Coupe du Maestrichtien à l'Oligocene inférieur dans le Flysch Externe de Cuediu (Bassin de la Bistrita). – In: Bombita, G. (ed.), *Micropaleontological Guide to the Mesozoic and Tertiary of the Romanian Carpathians*. Institute of Geology & Geophysics, Bucharest. 135-141.
- BROHI, I.A. (1994). – Late Cretaceous and Early Tertiary benthic foraminiferal biostratigraphy in Khuzdar District, Balachistan, Pakistan, with special reference to the influence of Tethyan Sea closure. – *Journal of Science of the Hiroshima University, Series C (Earth and Planetary Sciences)*, 10 (1), 25-86.
- BRÖNNIMANN, P. (1951). – Internal structure of *Cyclammina cancellata*. – *Journal of Paleontology*, 25, 765-761.
- BRÖNNIMANN, P. (1952). – Trinidad Paleocene and lower Eocene Globigerinidae. – *Bulletins of American Paleontology*, 34 (143), 1-34.
- BRÖNNIMANN, P. & WHITTAKER, J.E. (1987). – A revision of the foraminiferal genus *Adercotryma* Loeblich & Tappan, with a description of *A. wrighti* sp. nov. from British waters. – *Bulletin of the British Museum of Natural History (Zoology)*, 52, 19-28.
- BRÖNNIMANN, P. & WHITTAKER, J.E. (1988). – The Trochamminaceous test and the taxonomic criteria used in the classification of the superfamily Trochamminacea. – In: Proceedings of the Second Workshop on Agglutinated Foraminifera. Vienna 1986. *Abhandlungen der Geologischen Bundesanstalt*, 41, 13-40.
- BRÖNNIMANN, P. & ZANINETTI, L. (1984). – Agglutinated foraminifera mainly Trochamminacea from the Baía de Sepetiba, near Rio de Janeiro, Brazil. – *Revue de Paléobiologie*, 1, 63-115.

- BRÖNNIMANN, P., VAN DOVER, C.L. & WHITTAKER, J.E. (1989). – *Abyssotherma pacifica*, n.gen., n.sp., a recent remaneicid (Foraminiferida, Remaneicacea) from the East Pacific Rise. – *Micropaleontology*, 35, 142-149.
- BRÖNNIMANN, P., ZANINETTI, L. & WHITTAKER, J.E. (1983). – On the classification of the Trochamminacea (Foraminiferida). – *Journal of Foraminiferal Research*, 13 (3), 202-218.
- BROUWER, J. (1965). – Agglutinated foraminiferal faunas from some turbiditic sequences I, II. – *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen series B*, 68, 309-334.
- BUBÍK, M. (1995). – Vyskyt rodu *Eratidus* (Foraminifera) v Eocénu Moravského Flyše [Occurrence of the genus *Eratidus* in the Eocene of the Moravian flysch]. – Geological Research in Moravia and Silesia in the year 1994. Český geol. Ustav, Brno. 17-18. [in Czech].
- BUBÍK, M. (1995). – Cretaceous to Paleogene agglutinated foraminifera of the Bile Karpaty unit (West Carpathians, Czech Republic). – In: Kaminski, M.A., Geroch, S. & Gasinski, M.A. (eds), *Proceedings of the Fourth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 3, 71-116.
- BUTT, A. (1981). – Depositional environments of the Upper Cretaceous rocks in the northern part of the eastern Alps. – *Cushman Foundation for Foraminiferal Research Special Publication*, 20, 1-121.
- BYKOVA, N.K. (1939). – Foraminifery Verkhnomelovykh otlozhenii Ferganskoi doliny [Foraminifera of . – *Trudy Nauchno-Issledovatel'skogo Geologorazvedochnogo Instituta, series A*, 121, xx-xx.
- CARPENTER, W.R., PARKER, W.K. & JONES, T.R. (1862). – *Introduction to the study of Foraminifera*. Robert Hardwicke, London, 319 pp.
- CARTWRIGHT, N.G., GOODAY, A.J. & JONES, A.R. (1989). – The morphology, internal organization, and taxonomic position of *Rhizammina algaeformis* Brady, a large, agglutinated, deep-sea foraminifer. – *Journal of Foraminiferal Research*, 19 (2), 115-125.
- CHAMNEY, T.P. (1969). – Barremian Textulariina, Foraminifera from Lower Cretaceous beds, Mount Goodenough section, Aklavik Range, District of Mackenzie. – *Geological Survey of Canada Bulletin*, 185, 41 pp.
- CHAPMAN, F. (1904). – On some Cainozoic foraminifera from Brown's Creek, Otway Coast. – *Victoria Geol. Surv. Rec., Melbourne, Australia*, vol. 1, pt. 3, p. 229.
- CICHA, I. & ZAPETALOVÁ, I. (1963). – Wichtige Vertreter der Familie Lituolidae Reuss, 1861 (Foraminifera) aus dem Miozän der Westkarpaten. – *Sborník geologických Věd*, 1, 75-121.
- CICHA, I; RÖGL, F; RUPP, C; CTYROKA, J. (eds), (1998). – Oligocene-Miocene foraminifera of the Central Paratethys. – *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, 549, 1-325.
- CITA, M.B. & GRIGNINI, D. (1983). – Nature and origin of Late Neogene Mediterranean sapropels. – In: Schlanger, S.O. & Cita, M.B. (eds), *Nature and origin of Cretaceous carbon-rich sediments*. Academic Press, pp. 165-196.
- CHARNOCK, M.A. & JONES, R.W. (1990). – Agglutinated foraminifera from the Paleogene of the North Sea. – In: Hemleben, C., Kaminski, M.A., Kuhnt, W., and Scott, D.B. (eds), *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of Agglutinated Foraminifera*, NATO ASI Series, Kluwer Acad. Publ. 139-244.
- CHARNOCK, M.A. & JONES, R.W. (1997). – North Sea lituolid foraminifera with complex inner structures: taxonomy, stratigraphy and evolutionary relationships. – *Annales Societatis Geologorum Poloniae*, 67 (2-3), 183-196.
- CHEN, C.Y., LI, X.J. & CHEN, F. (1997). – Foraminiferal thanatocoensis of Eastern Pacific CC Area and its geological significance. – *Acta Micropaleontologica Sinica*, 14 (2), 215-223.
- COCCIONI, R., GALEOTTI, S. (1994). – K-T boundary extinction - geologically instantaneous or gradual event - evidence from deep-sea benthic foraminifera. – *Geology*, 22 (9), 779-782.
- COCCIONI, R., GALEOTTI, S. & GRAVILI, M. (1995). – Latest Albian - earliest Turonian deep-water agglutinated foraminifera in the Bottaccione section (Gubbio, Italy) – Biostratigraphic and palaeoecologic implications. *Revista Española de Paleontología, No. Homenaje al Dr. Guillermo Colom*, 135-152.
- COLE, F.E. (1981). – Taxonomic notes on the bathyal zone benthonic foraminiferal species off northeast Newfoundland. – *Bedford Institute of Oceanography Report*, BI-R-81-7. 121 pp.
- COLLINS, A.C. (1958). – Foraminifera. – Great Barrier Reef Expedition 1928-29, Scientific Reports, 6 (6), 335-437 + 5 pls.
- CORFIELD, R.M. & SHACKLETON, N.J. (1988). – Productivity change as a control on planktonic foraminiferal evolution after the Cretaceous/Tertiary boundary. – *Historical Biology*, 1, 323-343.

- CORFIELD, R.M., CARTLIDGE, J.E., PREMOLI SILVA, I. & HOUSLEY, R.A. (1991). – Oxygen and Carbon isotope stratigraphy of the Paleogene and Cretaceous limestones in the Bottaccione Gorge and the Contessa Highway sections, Umbria, Italy. – *Terra Nova*, 3, 414-432.
- CORLISS, B.H. (1985). – Microhabitats of benthic foraminifera within deep-sea sediments. – *Nature*, 314, 435-438.
- CORLISS, B.H. & CHEN, C. (1988). – Morphotype patterns of Norwegian Sea deep-sea benthic foraminifera and ecological implications. – *Geology*, 16, 716-719.
- CORLISS, B.H. & FOIS, E. (1991). – Morphotype analysis of deep-sea benthic foraminifera from the northwest Gulf of Mexico. – *PALAIOS*, 5, 589-605.
- CRESTA, S., MONECHI, S. & PARISI, G. (1989). – Mesozoic–Cenozoic stratigraphy in the Umbria-Marche area. – *Memorie Descrittive della Carta Geologica d'Italia*, 39, 182 pp.
- CUSHMAN, J.A. (1910). – A monograph on the foraminifera of the North Pacific Ocean; Part I - Astrorhizidae and Lituolidae. – *United States National Museum Bulletin*, 71, 134 pp.
- CUSHMAN, J.A. (1911). – A monograph on the foraminifera of the North Pacific Ocean; Part II - Textulariidae. – *United States National Museum Bulletin*, 71, 108 pp.
- CUSHMAN, J.A. (1917). – New species and varieties of foraminifera from the Philippines and adjacent waters. – *Proceedings of the US National Museum*, 51, 651-662.
- CUSHMAN, J.A. (1918). – The foraminifera of the Atlantic Ocean. Part 1. Astrorhizidae. – *United States National Museum Bulletin*, 104, 1-109+39 pl..
- CUSHMAN, J.A. (1920). – The foraminifera of the Atlantic Ocean. Part 2. Lituolidae. – *United States National Museum Bulletin*, 104, 1-109.
- CUSHMAN, J.A. (1921). – Foraminifera of the Philippine and adjacent seas - *United States National Museum Bulletin*, 100, 1-608.
- CUSHMAN, J.A. (1926). – The foraminifera of the Velasco Shale of the Tampico Embayment. – *American Association of Petroleum Geologists Bulletin*, 10, 581- 612.
- CUSHMAN, J.A. (1932). – *Textularia* and related forms from the Cretaceous. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 8, 86-96.
- CUSHMAN, J.A. (1934). The generic position of "*Cornuspira cretacea* Reuss". – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 10, 44-47.
- CUSHMAN, J.A. (1937a). – A monograph of the foraminiferal family Verneulinidae. – *Cushman Laboratory for Foraminiferal Research Special Publication*, 7, 1-157.
- CUSHMAN, J.A. (1937b). – A monograph of the foraminiferal family Valvulinidae. – *Cushman Laboratory for Foraminiferal Research Special Publication*, 8, 1-210.
- CUSHMAN, J.A. (1946). – Upper Cretaceous foraminifera of the Gulf Coastal region of the United States and adjacent areas. – *U.S. Geological Survey Professional Paper*, 206, 1-241.
- CUSHMAN, J.A. (1947). – A foraminiferal fauna from the Santa Anita Formation of Venezuela. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 23, 1-18.
- CUSHMAN, J.A. (1947). – A foraminiferal fauna from the Upper Cretaceous Arkadelphia Marl of Arkansas. – *U.S. Geological Survey Professional Paper*, 221-A, 17 pp.
- CUSHMAN, J.A. & BARKSDALE, J.D. (1930). – Eocene foraminifera from Martinez, California. – *Contributions from the Department of Geology of Stanford University*, 1 (2), 55-73.
- CUSHMAN, J.A. & BERMÚDEZ, P.J. (1937). – Further new species of foraminifera from the Eocene of Cuba. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 13, 1-29.
- CUSHMAN, J.A. & CAMPBELL, A.S. (1934). – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 10, 70-71.
- CUSHMAN, J.A. & GOUDKOV, P.P. (1944). – Some foraminifera from the Upper Cretaceous of California. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 20, 53-64.
- CUSHMAN, J.A. & JARVIS, P.W. (1928). – Cretaceous foraminifera from Trinidad. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 4, 85-103.
- CUSHMAN, J.A. & JARVIS, P.W. (1932). – Upper Cretaceous foraminifera from Trinidad. – *Proceedings of the United States National Museum*, 80, article 14, 1-60.
- CUSHMAN, J.A. & McMASTERS, J.H. (1936). – Middle Eocene foraminifera from the Llajas Formation, Ventura County, California. – *Journal of Paleontology*, 10 (6), 497-517.
- CUSHMAN, J.A. & RENZ, H.H. (1946). – The foraminiferal fauna of the Lizard Springs Formation of Trinidad, British West Indies. – *Cushman Laboratory for Foraminiferal Research Special Publication*, 18, 1-48.

- CUSHMAN, J.A. & RENZ, H.H. (1947). – The foraminiferal fauna of the Oligocene Ste. Croix Formation of Trinidad, B.W.I. – *Cushman Laboratory for Foraminiferal Research Special Publication*, 22, 46 pp.
- CUSHMAN, J.A. & RENZ, H.H. (1947). – Further notes on the Cretaceous foraminifera of Trinidad. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 23, 31-51.
- CUSHMAN, J.A. & RENZ, H.H. (1948). – Eocene foraminifera of the Navet and Hospital Hill formations of Trinidad, B.W.I. – *Cushman Laboratory for Foraminiferal Research Special Publication*, 24, xx-xx.
- CUSHMAN, J.A. & SIEGFUS, S.S. (1939). – Some new and interesting foraminifera from the Kreyenhagen Shale of California. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 15, 23-33.
- CUSHMAN, J.A. & SIEGFUS, S.S. (1942). – Foraminifera from the type area of the Kreyenhagen Shale of California. – *Transactions of the San Diego Society of Natural History*, 9, (34), 385-426.
- CUSHMAN, J.A. & STAINFORTH, R.M. (1945). – The foraminifera of the Cipero marl formation of Trinidad, British West Indies. – *Cushman Laboratory for Foraminiferal Research Special Publication*, 14, 91 pp.
- CUSHMAN, J.A. & WATERS, J.A. (1927). – Some arenaceous foraminifera from the Upper Cretaceous of Texas. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 2, 81-85.
- DAILEY, D.H. (1983). – Late Cretaceous and Paleocene benthic foraminifera from Deep Sea Drilling Project Site 516, Rio Grande Rise, western South Atlantic. – *Initial Reports of the Deep Sea Drilling Project*, 72, 757-782.
- DARAKCHIEVA, S. (1999). – Small Tertiary foraminifers from Bulgaria. Atlas. Sophia, Bulgaria, 92 pp.
- DAVIS, R. (2001). – *Benthic foraminifera at the Cretaceous/Tertiary boundary in Bottaccione Gorge, Italy: rapid or gradual faunal change?* – Unpublished M.Sc. report, Department of Geological Sciences, University College London, 80 pp.
- DE KLASZ, I. & DE KLASZ, S. (1990). – Danian deep water (bathyal) agglutinated Foraminifera from Bavaria and their comparison with approximately coeval agglutinated assemblages from Senegal and Trinidad. – In: Hemleben, C., Kaminski, M.A., Kuhnt, W. & Scott, D.B. (eds), *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of Agglutinated Foraminifera*. NATO ASI Series C327, 387-431.
- DELACA, T.E., 1986. – The morphology and ecology of *Astrammia rara*. – *Journal of Foraminiferal Research*, 16, 216-223.
- DE RIJK, S., TROELSTRA, S.R. & ROHLING, E.J. (1999). – Benthic foraminiferal distribution in the Mediterranean Sea. – *Journal of Foraminiferal Research*, 29 (2), 93-103.
- DE RIJK, S., JORISSON, F.J., ROHLING, E.J. & TROELSTRA, S.R. (2000). – Organic flux control on bathymetric zonation of Mediterranean benthic foraminifera. – *Marine Micropaleontology*, 40, 151-166
- DIAZ DE GOMERO, M.L. (1977). – Estratigrafía y micropaleontología del Oligoceno y Mioceno Inferior del Centro de la Cuenca de Falcón, Venezuela. – *GEOS* (escuela de Geología y Minas, UCV), 22, 3-60.
- DIAZ DE GOMERO, M.L. (1985). – Micropaleontología de la formación Agua Salada, Falcon Nororiental. In: Espejo, A. et al. (eds), *Estratigrafía, paleontología y sedimentación. IV Congreso Geológico Venezolano*. Sociedad Venezolana de Geólogos, 1, 384-453.
- DICKENS, G.R. (2000). – Methane oxidation during the late Paleocene thermal maximum. – *Bulletin de la Société Géologique de France*, 171, 37-49.
- DIDON, J., DURAND-DELGA, M. & KORNPROBST, J. (1973). – Homologies géologiques entre les deux rives du détroit de Gibraltar. – *Bulletin de la Société géologique de France* (7), 15, 77-105.
- D'IORIO, M.A. (1986). – Integration of foraminiferal and dinoflagellate data sets in quantitative stratigraphy of the Grand Banks and Labrador Shelf. – *Bulletin of Canadian Petroleum Geology*, 34, 277-283.
- DIXON, J., DIETRICH, J.R., McNEIL, D.H., McINTYRE, D.J., SNOWDON, L.R., BROOKS, P. (1985). – Geology, biostratigraphy and organic geochemistry of Jurassic to Pleistocene strata, Beaufort-MacKenzie area, Northwest Canada. Course Notes. – Canadian Society of Petroleum Geologists, Calgary, Alberta. ?? pp.
- D'ORBIGNY, A. (1846). – *Die fossilen Foraminiferen des tertiären Beckens von Wien*. – Gide et Comp. Paris, 312 pp.
- DYLAŻANKA, M. (1923). – Warstwy inoceramowe z łomu w Szymbarku koło Gorlic [The Inoceramus beds from a quarry in Szymbark near Gorlice] – *Rocznik Polskiego Towarzystwa Geologicznego*, 1, 36-80.
- EARLAND, A. (1933). – Foraminifera. Part II. South Georgia. – *Discovery Reports*, 7, 29-138.
- EARLAND, A. (1934). – Foraminifera; Part III - the Falklands sector of the Antarctic (excluding South Georgia). – *Discovery Reports*, 10, 1-208.
- EARLAND, A. (1936). – Foraminifera. Part IV. Additional records from the Weddell Sea sector from material obtained by the S.Y. 'Scotia'. – *Discovery Reports*, 13, 1-76.

- EGGER, J.G. (1910). – Ostracoden und Foraminiferen des Eybrunner Kreidemergels in der Umgebung von Regensburg. – *Ber. Naturwiss. Ver. Regensburg*, 12, 1-48 + 6 pls.
- EICHER, D. (1966). – Foraminifera from the Cretaceous Carlile Shale of Colorado. – *Contributions from the Cushman Foundation for Foraminiferal Research*, 17, 16-31.
- ELDRETT, J.S., HARDING, I.C., FIRTH, J.V. & ROBERTS, A.P. (2004). – Magnetostratigraphic calibration of Eocene–Oligocene dinoflagellate cyst biostratigraphy from the Norwegian–Greenland Sea. – *Marine Geology*, 204, 91-127.
- EMBERSON, S.K. (2003). – *Benthic foraminiferal response to global warming in the Early Eocene: Deep-water agglutinated foraminifera in Contessa Valley, Gubbio, Italy*. – Unpublished M.Sc. report, Department of Geological Sciences, University College London, 122 pp.
- EMILIANI, C. (1954). – The Oligocene microfaunas of the central part of the northern Apennines. – *Paleontographica Italica*, 48, 77-184.
- EVANS, J.R. & KAMINSKI, M.A. (1998). – Pliocene and Pleistocene chronostratigraphy and paleoenvironment of the central Arctic Ocean, using deep-water agglutinated foraminifera. – *Micropaleontology*, 44, 109-130.
- FINLAY, H.J. (1940a). – New Zealand Foraminifera: Key species in stratigraphy - No. 3. – *Transactions and Proceedings of the Royal Society of New Zealand*, 69, 309-329.
- FINLAY, H.J. (1940b). – New Zealand Foraminifera: Key species in stratigraphy - No. 4. – *Transactions and Proceedings of the Royal Society of New Zealand*, 69, 356-387.
- FINLAY, H.J. (1946). – The microfaunas of the Oxford Chalk and Eyre River Beds. – *Transactions and Proceedings of the Royal Society of New Zealand*, 76, 237-245.
- FLINT, J.M. (1899). – Recent Foraminifera, a descriptive catalogue of specimens dredged by the U.S. Fish Commission steamer Albatross. – *U.S. National Museum Annual Report (1897)*, pp. 249-349.
- FRANKE, A. (1912). – Die Foraminiferen des Unter-Eocenes der Ziegelei Schwarzenbeck. – *Jahrbuch der Königlich Preussischen Geologischen Landesanstalt und Bergakademie zu Berlin*, 32, 106-111.
- FRANKE, A. (1925). – Die Foraminiferen der pommerschen Kreide. – *Greifswald Universität Geologische-Palaeontologische Institut, Abhandlungen*, 6, 1-96.
- FREIMAN, E.V. (1970). – Ob izmenchivosti i sistematike reofaksov iz Paleogena Zapadno-Sibirskoy Nizmennosti. – In: Fursenko, A.V. (ed), *Obshchie Voprosy izycheniya Mikrofauny Sibiri, Dal'nego Vostoka i drugikh raionov*. Trudy Instituta Geologii i Geofiziki, Akademiya Nauk SSSR, Sibirskoye Otdelenie, 71, 105-113 + 2 pls.
- FRIEDBERG, W. (1901). – Otwornice warstw inoceramowych okolic Rzeszowa i Dębicy [The Foraminifera from the Inoceramus beds near Rzeszów and Dębica]. – *Rozprawy Wydziału matematyczno-przyrodniczego, Akademia Umiejętności w Krakowie, seria 2*, 41, 601-668.
- FRIZZEL, D.L. (1943). – Upper Cretaceous foraminifera from northwestern Peru. – *Journal of Paleontology*, 17, 331-353.
- FRIZZEL, D.L. (1954). – Handbook of Cretaceous foraminifera of Texas. – *Bureau of Economic Geology, University of Texas, Report of Investigations*, 22, 1-232.
- FRIZZEL, D.L. & SCHWARTZ, E. (1950). – A new litioloid foraminiferal genus from the Cretaceous, with an emendation of *Cribrostomoides* Cushman. – *University of Missouri, Technical Series, Bulletin*, 76, 1-12.
- FUENMAYOR, A.N. (1989). – *Manual de Foraminiferos de la Cuenca de Maracaibo*. – Division de Operaciones de Producción, Gerencia de Ingeniería de Petróleo, Gerencia de Geología, Maracaibo Venezuela, 191 pp.
- FURSENKO, A.V., TROITSKAYA, T.S., LEVCHUK, L.K., NESTEROVA, O.N., POLOVOVA, T.P. & FURSENKO, K.B. (1979). – Foraminifera of the Far-Eastern Seas of the USSR. – *Trudy Instituta Geologii i Geofiziki, Akademiya Nauk SSSR, Sibirskoye Otdelenie*, 387, 397 pp.
- GALEA-ALVAREZ, F.A. (1985). – *Biostratigraphy and depositional environment of the Upper Cretaceous - Eocene Santa Anita Group (North Eastern Venezuela)*. – Ph.D. Thesis, Free University Press, Amsterdam. 115 pp.
- GALEOTTI, S., ANGORI, E., COCCIONI, R., FERRARI, G., GALBRUN, B., MONECHI, S., PREMOLI SILVA, I., SPEIJER, R. & TURI, B. (2000). – Integrated stratigraphy across the Paleocene/Eocene boundary in the Contessa Road section, Gubbio (Central Italy). – *Bulletin de la Société géologique de France*, 171 (3), 355-365.
- GALEOTTI, S., BELLAGAMBA, M., KAMINSKI, M.A. & MONTANARI, A. (2002). – Deep sea benthic foraminiferal recolonisation following a volcanoclastic event in the lower Campanian of the Scaglia Rossa Formation (Umbria-Marche Basin, Central Italy). – *Marine Micropaleontology*, 44, 57-76.

- GALEOTTI, S., KAMINSKI, M.A., COCCIONI, R. & SPEIJER, R.P. (2004). – High-resolution Deep-water agglutinated foraminiferal record across the Paleocene/Eocene transition in the Contessa Road Section (Central Italy). – In: Bubík, M. & Kaminski, M.A., (eds), *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 8, 83-103.
- GELLAI-NAGY, A. (1988). – Delineation of Hantken's foraminiferal species from the original collection. – *A Magyar Allami Földtani Intézet Évi Jelentése 1988-évről*, II Rész, 133-173.
- GEROCH, S. (1959). – Stratigraphic significance of arenaceous foraminifera in the Carpathian flysch. – *Palaontologische Zeitschrift*, 33, 113-122.
- GEROCH, S. (1960). – Microfaunal assemblages from the Cretaceous and Paleogene Silesian Unit in the Beskid Śląski Mts. (western Carpathians). – *Biuletyn Instytutu Geologicznego*, 153, 7-138. Warszawa.
- GEROCH, S. (1966). – Lower Cretaceous small foraminifera of the Silesian series, Polish Carpathians. – *Rocznik Polskiego Towarzystwa Geologicznego*, 36, 413-480.
- GEROCH, S. & GRADZIŃSKI, R. (1955). – Stratigrafia serii podśląskiej Żywieckiego okna tektonicznego. – *Rocznik Polskiego Towarzystwa Geologicznego*, 24, 3-62.
- GEROCH, S., JEDNOROWSKA, A., KSIĄŻKIEWICZ, M. & LISZKOWA, J. (1967). – Stratigraphy based upon microfauna in the western Polish Carpathians. – *Biuletyn Instytutu Geologicznego*, 211, 186-282.
- GEROCH, S. & NOWAK, W. (1984). – Proposal of Zonation for the late Tithonian - late Eocene, based upon arenaceous foraminifera from the outer Carpathians, Poland. – In: Oertli, H. (Ed.) *Benthos '83; 2nd International Symposium on Benthic Foraminifera Pau (France), April 11-15, 1983*. Elf Aquitaine, ESSO REP and TOTAL CFP, Pau & Bourdeaux, pp. 225-239.
- GEROCH, S. & KAMINSKI, M.A. (1992). – The morphology and systematics of *Nothia excelsa* (Grzybowski), a deep-water agglutinated foraminifer. – *Rocznik Polskiego Towarzystwa Geologicznego*, 62, 255-265.
- GEROCH, S. & KOSZARSKI, L. (1988). – Agglutinated foraminiferal stratigraphy of the Silesian Flysch Trough. – In: *Proceedings of the Second Workshop on Agglutinated Foraminifera*. Vienna 1986. *Abhandlungen der Geologischen Bundesanstalt*, 41, 73-80.
- GEROCH, S. & OLSZEWSKA, B. (1990). – The oldest assemblages of agglutinated foraminifera of the Polish Flysch Carpathians. – In: Hemleben, C. et al. (eds), *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of Agglutinated Foraminifera*. NATO ASI Series C-327, 525-538. Kluwer Acad. Publ.
- GIBSON, G.W. (1967). – Foraminifera and stratigraphy of the Tongaporutuan Stage in the Taranaki coastal and six other sections. – *Transactions of the Royal Society of New Zealand, Geology*, 1(5), 1-70.
- GILLMORE, G.K. (1991). – *Palaeogene foraminifera of the Central and southern North Sea and their biostratigraphical application*. – Unpublished Ph.D. Thesis, University College London, 406 pp + 23 pls.
- GIVEN, M.M. & WALL, J.H. (1971). – Microfauna from the Upper Cretaceous Bearpaw Formation of south-central Alberta. – *Bulletin of Canadian Petroleum Geology*, 19, 504-546.
- GLAESSNER, M.F. (1937). – Studien über Foraminiferen aus der Kreide und dem Tertiär des Kaukasus; 1. Die Foraminiferen der ältesten Tertiärschichten des Nordwest-Kaukas. – *Problems of Paleontology*, 2-3, 349-408. Moscow.
- GOËS, A. (1894). – A synopsis of the Arctic and Scandinavian Recent marine Foraminifera hitherto discovered. – *Kongl. Svenska Vetenskaps Akademiens Handlingar*, 25 (9), 3-127 + 25 pls.
- GOLL, R.M. (1989). – A synthesis of Norwegian Sea biostratigraphies: ODP Leg 104 on the Vøring Plateau. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 104, 777-826.
- GOODAY, A.J. & COOK, P.L. (1984). – An association between komokiacean foraminifera (Protozoa) and paludicelline ctenostomes (Bryozoa) from the abyssal northeast Atlantic. – *Journal of Natural History*, 18, 765-784.
- GOODAY, A.J. (1988). – A response by benthic foraminifera to the deposition of phytodetritus in the deep sea. – *Nature*, 332, 70-73.
- GOODAY, A.J. (1993). – Deep-sea benthic foraminiferal species which exploit phytodetritus: Characteristic features and controls on distribution. – *Marine Micropaleontology*, 22, 187-205.
- GOODAY, A.J. (1983). – Primitive foraminifera and xenophyophores in IOS epibenthic sledge samples from the northeast Atlantic. – *Institute of Ocean Sciences Report*, 156, 33 pp.
- GOODAY, A.J. & NOTT, J.A. (1982). – Intracellular barite crystals in two xenophyophores, *Aschemonella ramuliformis* and *Galatheammia* sp. (Protozoa: Rhizopoda) with comments on the taxonomy of *A. ramuliformis*. – *Journal of the Marine Biological Association of the United Kingdom*, 62, 595-605.

- GRADSTEIN, F.M. (1983). – Cenozoic Stratigraphy using agglutinated foraminifera. – Proceedings of the First Workshop on Agglutinated Foraminifera, Amsterdam, Sept. 1981. *Continental Shelf Institute Special Publication*, 108, 247-250.
- GRADSTEIN, F.M. & AGTERBERG, F.P. (1998). – Uncertainty in stratigraphic correlation. – *Norwegian Petroleum Society Special Publication*, 8, 9-29.
- GRADSTEIN, F.M. & AGTERBERG, F.P. (1982). – Models of Cenozoic Foraminiferal Stratigraphy--Northwestern Atlantic Margin. – In: Cubitt, J.M. & Reyment, R.A. (eds), *Quantitative Stratigraphic Correlation*. John Wiley & Sons.
- GRADSTEIN, F.M. & BÄCKSTRÖM, S.A. (1996). – Cainozoic biostratigraphy and palaeobathymetry, northern North Sea and Haltenbanken. – *Norsk Geologisk Tidsskrift*, 76, 3-32.
- GRADSTEIN, F.M. & BERGGREN, W.A. (1981). – Flysch-type agglutinated foraminifera and the Maestrichtian to Paleogene history of the Labrador and North Seas. – *Marine Micropaleontology*, 6, 211-268.
- GRADSTEIN, F.M. & KAMINSKI, M.A. (1989). – Taxonomy and biostratigraphy of new and emended species of Cenozoic deep-water agglutinated foraminifera from the Labrador and North Seas. – *Micropaleontology*, 35, 72-92.
- GRADSTEIN, F.M. & KAMINSKI, M.A. (1997). – New species of Paleogene deep-water agglutinated foraminifera from the North Sea and Norwegian Sea. – *Annales Societatis Geologorum Poloniae*, 67, 217-229.
- GRADSTEIN, F.M. & SRIVASTAVA, S.P. (1980). – Aspects of Cenozoic stratigraphy and paleoceanography of the Labrador Sea and Baffin Bay. – *Paleogeography Paleoclimatology Paleocology*, 30, 261-295.
- GRADSTEIN, F.M. & WILLIAMS, G.L. (1976). – Biostratigraphy of the Labrador Shelf, Part 1. – *Geological Survey of Canada Open File*, 349, 1-39.
- GRADSTEIN, F.M., KAMINSKI, M.A. & AGTERBERG, F.P. (1999). – Biostratigraphy and paleoceanography of the Cretaceous seaway between Norway and Greenland. – *Earth Science Reviews*, 46 (1-4), 27-98.
- GRADSTEIN, F.M., KAMINSKI, M.A. & BERGGREN, W.A. (1988). – Cenozoic foraminiferal biostratigraphy, Central North Sea. – In: Proceedings of the Second Workshop on Agglutinated Foraminifera. Vienna 1986. *Abhandlungen der Geologischen Bundesanstalt*, 41, 97-108.
- GRADSTEIN, F.M., KAMINSKI, M.A., BERGGREN, W.A. & D'IORIO, M.A. (1994). – Cenozoic biostratigraphy of the Central North Sea and Labrador Shelf. – *Micropaleontology* vol. 40 Supplement, 152 pp.
- GRADSTEIN, F.M., KRISTIANSEN, I.L., LOEMO, L. & KAMINSKI, M.A. (1992). – Cenozoic foraminiferal and dinocyst stratigraphy of the central North Sea. – *Micropaleontology*, 38, 101-137.
- GREEN, R.C., KAMINSKI, M.A. & SIKORA, P.J. (2004). – Miocene deep water agglutinated foraminifera from Viosca Knoll, offshore Louisiana (Gulf of Mexico). – In: Bubík, M. & Kaminski, M.A. (eds), *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 8, 119-144.
- GRIMM, K.I. (2002). – Foraminiferal zonation of early Oligocene deposits (Selztal Group, Latdorfian, Rupelian) in the Mainz Basin, Germany. – *Journal of Micropalaeontology*, 21, 67-74.
- GRÜN, W., LAUER, G., NIEDERMAYER, G. & SCHNABEL, W. (1964). – Die Kreide- Tertiär Grenze im Wienerwaldflysch bei Hochstrass/Niederösterreich. – *Verhandlungen der Geologischen Bundesanstalt*, 2, 226-283. Wien.
- GRÜN, W. (1969). – Flysch microfauna of the Hagenbach Valley (Northern Vienna Woods) Austria. – *Rocznik Polskiego Towarzystwa Geologicznego*, 39, 305-334.
- GRÜN, W. (1972). – Studien in der Unterkreide des Wienerwaldes. – *Jahrbuch der Geologischen Bundesanstalt*, 115, 2, 103 - 186.
- GRZYBOWSKI, J. (1896). – Otwornice czerwonych iłów z Wadowic. – *Rozprawy Wydziału Matematyczno-Przyrodniczego, Akademia Umiejętności w Krakowie, seria 2*, 30, 261-308.
- GRZYBOWSKI, J. (1898a). – Otwornice pokładów naftonosnych okolicy Krosna. – *Rozprawy Wydziału Matematyczno-Przyrodniczego, Akademia Umiejętności w Krakowie, seria 2*, 33, 257-305.
- GRZYBOWSKI, J. (1898b). – Mikroskopowe badania namulów wiertniczych z kopalń naftowych. 1. Pas potocki i okolice Krosna. 2. Uwagi ogólne. – *Kosmos (Lwów)*, 22, 393-439.
- GRZYBOWSKI, J. (1901). – Otwornice warstw inoceramowych okolicy Gorlic. – *Rozprawy Wydziału Matematyczno-Przyrodniczego, Akademia Umiejętności w Krakowie, seria 2*, 41, 219-286.
- GUZIK, K & POŻARYSKI, W. (1949). – Fald Biecza. – *Biuletyn Państwowego Instytutu Geologicznego*, 53, 33 pp.

- HAGN, H. (1960). – Die stratigraphischen, paläogeographischen und tektonischen Beziehungen zwischen Molasse und Helvetikum im östlichen Oberbayern. – *Geologica Bavarica*, 44, 1 - 208.
- HAIG, D.W. (1982). – Deep sea foraminiferids from Paleocene sediments, Port Moresby, Papua New Guinea. – *Journal of Foraminiferal Research*, 12, 287-297.
- HANAGATA, S., MOTOYAMA, I. & MIWA, M. (2001). – Geologic ages of the last occurrence of *Spirosigmoinella compressa* and the first occurrence of *Miliammina echigoensis* (benthic foraminifera), and their paleoceanographic implications – response to the latest Miocene–earliest Pliocene sea level changes. – *Journal of the Geological Society of Japan*, 107 (2), 101-116.
- HANTKEN, M. von (1868). – A kis-czelli tályag foraminiferái. – *Magyar Foldtani Társulat Munkálatai, Pest*, 4, 75-96.
- HANTKEN, M. von (1875). – Die Fauna der Clavulina szaboi-Schichten; Theil I Foraminiferen. – *Mittheilungen aus dem Jahrbuche der Koniglich-Ungarische Geologische Anstalt, Budapest*, 4 (1), 1-93 + 13 pls.
- HANZLÍKOVÁ, E. (1953). – Mikropaleontologicko-stratigrafické zhodnocení vrstvy Žukov NP15. – *Sborník Ústředního Ústavu Geologického, oddíl paleontologický*, 20, 1-83 + 13 pls.
- HANZLÍKOVÁ, E. (1955). – Príspevek k mikrobiostratigrafii magurského flyše v povodí Oravy (List spec. mapy Tvrdošín 4263). – *Zpr. geol. Vyzk. v Roce 1954*, 22 - 25.
- HANZLÍKOVÁ, E. (1958). – Předběžná zpráva o mikrobiostratigrafických poměrech Ľubovnianské vrchoviny – *Vestník Ústředního Ústavu Geologického*, 33, 429-440.
- HANZLÍKOVÁ, E. (1965). – Stratigraphie der Kreide und des Palaogens der flyschzone der Westkarpaten. – *Geologický Sborník*, 16, 33-64. Bratislava.
- HANZLÍKOVÁ, E. (1966). – Die Foraminiferen der Lhoty-Schichten. – *Acta Musei Moraviae, - Scientiae Naturales*, 51, 95-132. Brno.
- HANZLÍKOVÁ, E. (1969). – The foraminifera of the Frydek Formation (Senonian). – *Sborník geologických Věd, Paleontologie*, 11, 7 - 84.
- HANZLÍKOVÁ, E. (1972a). – Mikropaläontologische Zonenteilung und Stratigraphie der Istebna-Schichten und ihres unmittelbaren Hangenden. – *Vestník Ústředního Ústavu Geologického*, 47, 2, 69 - 77.
- HANZLÍKOVÁ, E. (1972b). – Carpathian Upper Cretaceous foraminifera from Moravia (Turonian-Maestrichtian). – *Rozpravy Ústředního Ústavu Geologického*, 39, 5-160.
- HANZLÍKOVÁ, E. (1973). – Foraminifera of the variegated Godula Member in Moravia (Cenomanian-Turonian). – *Sborník geologických Věd, Paleontologie*, 15, 119-184. Praha.
- HANZLÍKOVÁ, E. (1983). – Paleogene stratigraphy and foraminifera of the Outer Flysch Belt. – *Miscellanea Micropaleontologica; Knihovnicka Zemního Plynu a Nafty*, 4, 43-71.
- HANZLÍKOVÁ E., PÍCHA F. & ČICHA I. (1963). – Geologie karpatských pokryvných útvarů na vrtném poli Chlebovice - Rychaltice (Beskydy). – *Sborník Ústředního Ústavu Geologického, oddíl geologický*, 28, 435 - 455.
- HANZLÍKOVÁ, E. & PESL V. (1964) – Príspevek k poznání faciálního vývoje spodního oddílu paleogénu račanské série magurského flyše. – *Vestník Ústředního Ústavu Geologického*, 39, 6, 419 - 428.
- HARLOFF, J. & MACKENSEN, A. (1997). – Recent benthic foraminiferal associations and ecology of the Scotia Sea and Argentine Basin. – *Marine Micropaleontology*, 31, 1-29.
- HARRIS, R.W. & JOBE, B.I. (1951). – *Microfauna of basal midway outcrops near Hope, Arkansas*. – Transcript Press, Norman OK.
- HART, M.B. & BIGG, P.J. (1983). – Arenaceous foraminifera from the late Cretaceous "Anoxic Events" in northwest Europe. – Proceedings of the First Workshop Arenaceous Foraminifera. *Continental Shelf Institute Special Publication*, 108, 89-90.
- HEDBERG, H. (1937). – Foraminifera of the middle Tertiary Carapita formation of northeastern Venezuela. – *Journal of Paleontology*, 11, 661-697.
- HEDLEY, R.H. (1963). – Cement and iron in the arenaceous foraminifera. – *Micropaleontology*, 9, 433-441.
- HERON-ALLEN, E. (1915). – A short statement upon the theory, and the phenomena of purpose and intelligence exhibited by the Protozoa, as illustrated by selection and behaviour in the Foraminifera. – *Journal of the Royal Microscopical Society*, 1915, 547-557.
- HERON-ALLEN, E. & EARLAND, A. (1913). – On some foraminifera from the North Sea, etc., dredged by the Fisheries Cruiser 'Goldseeker' (International North Sea Investigations – Scotland). II. On the distribution of *Saccammina sphaerica* (M.Sars) and *Psammosphaera fusca* (Schultze) in the North Sea: particularly with reference to the suggested identity of the two species. – *Journal of the Royal Microscopical Society*, v. 1913, 1-26.
- HEMLEBEN, C. & TROESTER, J. (1984). – Campanian-Maestrichtian deep-water foraminifers from Hole 543A, Deep Sea Drilling Project. – *Initial Reports of the Deep Sea Drilling Project*, 78A, 509-532.

- HEMLEBEN, C. & KAMINSKI, M.A. (1990). – Agglutinated foraminifera: an introduction. – In: Hemleben, C., Kaminski, M.A., Kuhnt, W. & Scott, D.B. (eds), *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of agglutinated foraminifera*. NATO ASI C, 327, 3-11. Kluwer Academic Press.
- HESS, S. (1998). – Verteilungsmuster rezenter benthischer Foraminiferen im Südchinesischen Meer. – *Berichte – Reports, Geologisch-Paläontologisches Institut und Museum, Christian-Albrechts-Universität Kiel*, 91, 173 pp.
- HESS, S. & KUHN, W. (1996). – Deep-sea benthic foraminiferal recolonization of the 1991 Mt. Pinatubo ash layer in the South China Sea. – *Marine Micropaleontology*, 28, 171-197.
- HESS, S., KUHN, W., SPIVEY, B., KAMINSKI, M.A. & WHITTAKER, J.E. (2000). – Ash grains of the 1991 Mt. Pinatubo eruption as a tracer in Rose Bengal stained deep sea agglutinated foraminifera: How old is Freddy? – In: Hart, M.B., Kaminski, M.A. & Smart, C.W. (eds), *Proceedings of the Fifth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 7, 126.
- HESS, S., KUHN, W., HILL, S., KAMINSKI, M.A., HOLBOURN, A. & DE LEON, M. (2001). – Monitoring the recolonisation of the Mt. Pinatubo 1991 ash layer by benthic foraminifera. – *Marine Micropaleontology*, 43, 119-142.
- HESSE, R. & BUTT, A. (1976). – Paleobathymetry of Cretaceous turbidite basins of the east Alps relative to the calcite compensation level. – *Journal of Geology*, 84, 505-533.
- HILLEBRANDT, A. von (1962). – Das Paleozän und seine Foraminiferen-fauna im Becken von Reichenhall und Salzburg. – *Abhandlungen, Bayerische Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, 108, 9-180. Munich.
- HILLEBRANDT, A. von (1965). – Foraminiferen-Stratigraphie im Alttertiär von Zumaya (Provinz Guipuzcoa, N.W.-Spanien) und ein Vergleich mit anderen Tethys-Gebieten. – *Abhandlungen, Bayerische Akademie Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse*, Heft 123. 62 pp.
- HILTERMANN, H.H. (1943). – Zur Stratigraphie und Mikrofossilführung der Mittelkarpaten. – *Erdöl und Kohle*, 39, 33/34, 745 - 755.
- HILTERMANN, H.H. (1968). – Neuere paleontologische Daten zum Flysch-Problem. – *Erdöl und Erdgas*, 84, 151-157.
- HILTERMANN, H.H. (1972). – Zur Morphologie der Benthos-Foraminifere *Spiroplectammina spectabilis* (Grzybowski). – *Geologisches Jahrbuch*, Reihe A4, 43-61.
- HILTERMANN, H.H. (1973). – Zur Sociologie von Agglutinierenden Foraminiferen polarer skandinavischer und aquatorialer Meere. – *Geologische Jahrbuch*, Reihe A6, 101-120.
- HILTERMANN, H.H. (1974). – *Rzehakina epigona* und unterarten dieser Foraminifere. – *Palaontologische Zeitschrift*, 48, 36-56.
- HILTERMANN, H.H. (1975). – Zur stratigraphie und Mikrofossil-führung der Mittelkarpaten – Münstersche Forschungen zur Geologie und Paläontologie, 35, 1-50.
- HOFKER, J. (1959). – Les foraminifères du Crétacé Supérieur du Cotentin, – 84 Congres des Societes Savantes, Dijon, pp. 369-397.
- HOFKER, J. (1966). – Maestrichtian, Danian and Paleocene foraminifera. The foraminifera of the type Maestrichtian in south Limburg, Netherlands, together with the foraminifera of the underlying Gulpen Chalk and the overlying calcareous sediments. The foraminifera of the Danske chalk and the overlying greensands as found in Denmark. – *Paleontographica*, 40, 1-376. Stuttgart.
- HOFKER, J. (1972). – *Primitive agglutinated foraminifera*. – E.J. Brill, Leiden. 95 pp.
- HOFKER, J. (1976) Further studies on Caribbean Foraminifera. – *Studies on the Fauna of Curacao and other Caribbean Islands*, 49 (162), 1-256.
- HÖGLUND, H. (1947). – Foraminifera in the Gullmar Fjord and the Skagerak. – *Zoologiska Bidrag Från Uppsala*, 26, 4-328.
- HOLBOURN, A.E.L. & KAMINSKI, M.A. (1997). – Lower Cretaceous benthic foraminifera of the Indian Ocean. – *Grzybowski Foundation Special Publication*, 4, 172pp.
- HOLZKNECHT, M. (1988). – Aglutinující foraminifery středního Eocénu zlínského souvrství z vrtu v okolí Vyšné Radváně (V. Slovensko, ČSSR) [Aglutinierende Foraminiferen des Mitteleozäns des Zlín-Schichtenkomplexes aus den Bohrungen der Umgebung von Vyšná Radván (Ostslowakei, Tschechoslowakei)]. – *Knihovnička Zemního plynu a nafty, Miscellanea Micropalaeontologica*, 3, 73-95.
- HOLZKNECHT, M. & KRHOVSKY, J. (1987). – Paleocenní až spodnoeocenní foraminifery nesvačilkého souvrství opěrné vrtby Nesvačilka-1 (autochton vnújsich Západních Karpat na j. Moravě) [Paleocene to early Eocene foraminifera of the Nesvačilka Formation from Nesvačilka-1 borehole (the autochthon of the external western Carpathians, southern Moravia, Czechoslovakia)]. – *Knihovnička Zemního plynu a nafty, Miscellanea Micropalaeontologica*, 2(2), 127-215.

- HOMOLA, V. & HANZLIKOVÁ, E. (1955) – Biostratigrafické, tektonické, a lithologické studie na Těšínsku. – *Sbornik Ústředního Ústavu Geologického*, 21, 317-498.
- HORNIBROOK, N. (1961). – Tertiary foraminifera from the Oamaru District (N.Z.). – *New Zealand Geological Survey Bulletin*, 34, 191 pp.
- HORNIBROOK, N., BRAZIER, R.C. & STRONG, C.P. (1989). – Manual of Permian to Pleistocene foraminiferal biostratigraphy. – *New Zealand Geological Survey Paleontological Bulletin*, 56, 175 pp.
- HORNIBROOK, N. & HAY, R.F. (1978). – Late Cretaceous agglutinated foraminifera from sediments interbedded with the Tangihua Volcanics, Northland, New Zealand. – *BMR Bulletin*, 192, 67-72.
- HORVÁTH, M. (2002). – Data to revision and distribution of small Foraminifera species described by Hantken (1868, 1875). Part 1. Textulariidae and Miliolidae. – *Fragmenta Palaeontologica Hungarica*, 20, 25-42.
- HOTTINGER, L., HALICZ, E. & REISS, Z. (1990). – Wall texture of *Spirorutilus*. – *Journal of Foraminiferal Research*, 20, 65-70.
- HULL, D.M., OSTERMAN, L.E. & THIEDE, J. (1996). – Biostratigraphic synthesis of Leg 151, North Atlantic – Arctic Gateways. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 151, 627-644.
- HULSBOS, R.E., KROON, D., JANSEN, H.S.M. & VAN HINTE, J.E. (1989). – Lower Eocene benthic foraminifera and paleoenvironment of the outer Voring Plateau, Norwegian Sea (DSDP Site 338). – *Micropaleontology*, 35, 256-273.
- HUSS, F. (1966). – Otwornice aglutynujące serii podślaskiej jednostki roponośnej Węglowki (Polskie Karpaty Fliszowe) [Agglutinated foraminifera of the oil-bearing subsilesian series in Weglowka (Polish Fylsch Carpathians)]. – *Prace Geologiczne, Polska Akademia Nauk*, 34, 7-76.
- ISRAELSKY, M.C. (1951). – Foraminifera of the Lodo Formation Central California. – *U.S. Geological Survey Professional Paper*, 240-A, 29 pp.
- ION-SANDULESCU, J. (1976). – Microbiostratigraphie, associations et zones a foraminifères du Crétacé du flysch externe des Carpates Orientales (Roumanie). – *Revista Española de Micropaleontologia*, 7 (1), 99-111.
- ION, J. (1995). – Stratigraphy of the late Maastrichtian-Paleocene deposits of the Vrancea Halfwindow (Marginal Folds Nappe, East Carpathians - Romania) according to Foraminifera. – *Romanian Journal of Paleontology*, 76, 21-30.
- JEDLITSCHKA, H. (1935) - Beitrag zur Kenntnis der Mikrofauna der subbeskidischen Schichten. – *Mitt. naturwiss. Ver. (Troppau)*, 40, 27, 31 - 48.
- JEDNOROWSKA, A. (1968). – Zespoły otwornicowe w zewnętrznych strefach jednostki magurskiej Karpat i ich znaczenie stratigraficzne [Foraminiferal assemblages in the external zone of the Magura Unit of the Carpathians and their stratigraphic significance]. – *Prace Geologiczne, Polska Akademia Nauk*, 50, 7-89.
- JEDNOROWSKA, A. (1975). – Zespoły małych otwornic w paleocenie Polskich Karpat zachodnich [Smaller foraminifera assemblages in the Paleocene of the Polish Western Carpathians]. – *Studia Geologica Polonica*, 47, 7-103.
- JOHN, A.W.G. (1987). – The regular occurrence of *Reophax scotii* Chaster, a benthic foraminiferan in plankton samples from the North Sea. – *Journal of Micropalaeontology*, 6 (2), 61-63.
- JONASSON, K.E., SCHRÖDER-ADAMS, C.J. & PATTERSON, RT. (1995). – Benthic foraminiferal distribution at Middle Valley, Juan De Fuca Ridge, A Northeast Pacific hydrothermal venting site. – *Marine Micropaleontology*, 25, 151-167.
- JONES, G.D. (1988). – A paleoecological model of late Paleocene "flysch-type" agglutinated foraminifera using the paleoslope transect approach, Viking Graben, North Sea. – *In: Proceedings of the Second Workshop on Agglutinated Foraminifera. Vienna 1986. Abhandlungen der Geologischen Bundesanstalt*, 41, 143-154.
- JONES, R.W. (1994). – *The Challenger Foraminifera*. – Oxford University Press, 149+vi pp + 117 pl.
- JONES, R.W. & CHARNOCK, M.A. (1985). – "Morphogroups" of agglutinating foraminifera. Their life positions and feeding habits and potential applicability in (paleo)ecological studies. – *Revue de Paleobiologie*, 4, 311-320.
- JONES, R.W., BENDER, H., CHARNOCK, M.A., KAMINSKI, M.A. & WHITTAKER, J.E. (1993). – Emendation of the foraminiferal genus *Cribrostomoides* Cushman, 1910, and its taxonomic implications. – *Journal of Micropalaeontology*, 12 (2), 181-193.
- JONES, T.R. & PARKER, W.K. (1860). – On the Rhizopodal fauna of the Mediterranean compared with that of the Italian and some other Tertiary deposits. – *Quarterly Journal of the Geological Society of London*, 16, 292-307.

- JUMARS, P.A. MAYER, L.A., DEMINO, J.W., BAROSS, J.A. & WHEATCROFT, R.A. (1990). – Deep-sea deposit feeding strategies suggested by environmental and feeding constraints. – *Philosophical Transactions of the Royal Society of London*, A331, 85-101
- JURKIEWICZ, H. (1960). – Otwornice z łupków czarnorzeczki wschodniej części jednostki śląskiej [Foraminifera from the Czarnorzeczka shales of the eastern part of the Silesian Unit]. – *Rocznik Polskiego Towarzystwa Geologicznego*, 30, 333-344.
- JURKIEWICZ, H. (1967). – Foraminifera in the sub-Menilitic Paleogene of the Polish Middle Carpathians. – *Biuletyn Instytutu Geologicznego*, 210, 5-116. Warszawa.
- KAIHO, K. (1984). – Paleogene foraminifera from Hokkaido, Japan. Part 1. Lithostratigraphy and biostratigraphy including description of new species. – *Science Reports of the Tohoku University, Sendai, Second Series (Geology)*, 54 (2), 95-139.
- KAIHO, K. (1984). – Paleogene foraminifera from Hokkaido, Japan. Part 2. Correlation of the Paleogene System in Hokkaido and systematic paleontology. – *Science Reports of the Tohoku University, Sendai, Second Series (Geology)*, 55 (1), 1-74.
- KAIHO, K. (1991). – Global changes of Paleogene aerobic/anaerobic benthic foraminifera and deep-sea circulation. – *Palaeogeography Palaeoclimatology Palaeoecology*, 83, 65-85.
- KAIHO, K. (1992). – A low extinction rate of intermediate-water benthic foraminifera at the Cretaceous/Tertiary boundary. – *Marine Micropaleontology*, 18, 229-259.
- KAIHO, K. & HASEGAWA, T. (1994). – End-Cenomanian benthic foraminiferal extinctions and oceanic dysoxic events in the northern Pacific Ocean. – *Paleogeography Paleoclimatology Paleoecology*, 111, 29-43.
- KAIHO, K., FUJIWARA, O. & MOTOYAMA, I. (1993). – Mid-Cretaceous faunal turnover of intermediate-water benthic foraminifera in the northeastern Pacific Ocean margin. – *Marine Micropaleontology*, 23, 13-49.
- KAIHO, K., ARINOBU, T., ISHIWATARI, R., MORGANS, H.E.G., OKADA, H., TAKEDA, N., TAZAKI, K., ZHOU, G., KAJIWARA, Y., MATSUMOTO, R., HIRAI, A., NIITSUMA, N. & WADA, H. (1996). – Latest Paleocene benthic foraminiferal extinction and environmental changes at Tawanui, New Zealand. – *Paleoceanography*, 11, 447-465.
- KALISHEVICH, T.G., ZAKLINSKAYA, E.D. & SEROVA, M.Ya. (1981). – *Razvitie organicheskogo Mira Tikhookeanskogo Poyasa na Rubezhe Mezozoya i Kainozoya: Foraminifery, Mollyuski i Palinoflora*. – Izdatelstvo "Nauka", 144 pp.
- KAMINSKI, M.A. (1983). – Taxonomic notes on the abyssal agglutinated benthic foraminifera of the HEBBLE Area (lower Nova Scotian Continental Rise). – *Woods Hole Oceanographic Institution Technical Report*, WHOI-83-35, 49 pp.
- KAMINSKI, M.A. (1984). – Shape variation in *Spiroplectamina spectabilis* (Grzybowski). – *Acta Paleontologica Polonica*, 29, 29-49.
- KAMINSKI, M.A. (1985). – Evidence for control of abyssal agglutinated foraminiferal community structure by substrate disturbance: Results from the HEBBLE Area. – *Marine Geology*, 66, 113-131.
- KAMINSKI, M.A. (1988). – *Cenozoic deep-water agglutinated foraminifera in the North Atlantic*. – Ph.D. Thesis, WHOI/MIT Joint Program in Oceanography. WHOI 88-3, pp. 262.
- KAMINSKI, M.A. (2004). – The Year 2000 Classification of the Agglutinated Foraminifera. – In: Bubík, M. & Kaminski, M.A., (eds), *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 8, 237-255.
- KAMINSKI, M.A. & AUSTIN, W.E.N. (1999). – Oligocene Deep Water Agglutinated Foraminifera at Site 985, Norwegian Basin, southern Norwegian Sea. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 162, 169-177.
- KAMINSKI, M.A. & CRESPO DE CABRERA, S. (1999). – A new species of primitive *Reticulophragmium* (Foraminifera) from the Paleocene Vidoño Formation of northeastern Venezuela. – *Annales Societatis Geologorum Poloniae*, 69, 189-193.
- KAMINSKI, M.A. & FILIPESCU, S. (2000). – *Praesphaerammina*, a new genus of Cenozoic deep-water agglutinated foraminifera from the Carpathian flysch deposits. – *Micropaleontology*, 46, 353-359.
- KAMINSKI, M.A. & GEROCH, S. (1987). – Two new species of *Phenacophragma* from the Paleogene of Trinidad and Poland. – *Micropaleontology*, 33, 185-188.
- KAMINSKI, M.A. & GEROCH, S. (1992). – *Trochamminoides grzybowskii*, nom.nov., a new name for *Trochammina elegans* Grzybowski, 1898 (Foraminiferida). – *Journal of Micropalaeontology*, 11, 137.
- KAMINSKI, M.A. & GEROCH, S. (1993). – A revision of foraminiferal species in the Grzybowski Collection. – In: Kaminski, M.A., Geroch, S. & Kaminski, D. (eds), *The Origins of Applied Micropaleontology: The School of Jozef Grzybowski*. Grzybowski Foundation Special Publication no. 1. Alden Press, Oxford, 239-323.

- KAMINSKI, M.A. & GEROCH, S. (1997). – *Psamminopelta gradsteini* n.sp., a new species of Paleogene deep-water agglutinated foraminifera from the northern North Atlantic and Polish Outer Carpathians. *In: Hass, H.C. & Kaminski, M.A. (eds), Contributions to the Micropaleontology & Paleoceanography of the Northern North Atlantic*. Grzybowski Foundation Special Publication, 5, 249-252.
- KAMINSKI, M.A. & GRADSTEIN, F.M. (1987). – Paleobiogeography of Paleogene flysch-type foraminiferal assemblages in the North Atlantic. – Gulf Coast Section/SEPM Foundation 8th Annual Research Conference, Selected papers and illustrated abstracts, pp. 85-89.
- KAMINSKI, M.A. & HUANG, Z. (1991). – Biostratigraphy of deep-water agglutinated foraminifera at Site 767 (Celebes Sea). – *Proceedings of the Ocean Drilling Program, Scientific Results*, 124, 171-180.
- KAMINSKI, M.A. & KUHN, W. (1991). – Depth-related shape variation in *Ammobaculites agglutinans* (d'Orbigny). – *Annales Societatis Geologorum Poloniae*, 61, 221-230.
- KAMINSKI, M.A. & SCHRÖDER, C.J. (1987). – Environmental analysis of deep-sea agglutinated foraminifera: Can we distinguish tranquil from disturbed environments? - Gulf Coast Section/SEPM Foundation 8th Annual Research Conference, Selected papers and illustrated abstracts, pp. 90-93.
- KAMINSKI, M.A. & WETZEL, A. (2004a). – Sediment disturbance caused by a suspension-feeding tubular agglutinated foraminifera. – *In: Bubik, M. & Kaminski, M.A., (eds), Proceedings of the Sixth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 8, 294.
- KAMINSKI, M.A. & WETZEL, A. (2004b). – A tubular protozoan predator: A burrow selectively filled with tubular agglutinated protozoans (Xenophyophorea, Foraminifera) in the abyssal South China Sea. – *In: Bubik, M. & Kaminski, M.A., (eds), Proceedings of the Sixth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 8, 287-293.
- KAMINSKI, M.A., BOERSMA, A., TYSZKA, J. & HOLBOURN, A.E.L. (1995). – Response of deep-water agglutinated foraminifera to dysoxic conditions in the California borderland basins. – *In: Kaminski, M.A., Geroch, S. & Gasinski, M.A. (eds), Proceedings of the Fourth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 3, 131-140.
- KAMINSKI, M.A., GRADSTEIN, F.M., BERGGREN, W.A., GEROCH, S. & BECKMANN, J.P. (1988). – Flysch-type agglutinated foraminiferal assemblages from Trinidad: Taxonomy, Stratigraphy and Paleobathymetry. – *In: Proceedings of the Second Workshop on Agglutinated Foraminifera, Vienna 1986. Abhandlungen der Geologischen Bundesanstalt*, 41, 155-228.
- KAMINSKI, M.A., GRADSTEIN, F.M., SCOTT, D.B. & MacKINNON K.D. (1989). – Neogene benthic foraminiferal stratigraphy and deep water history of Sites 645, 646 and 647, Baffin Bay and Labrador Sea. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 105, 705-730.
- KAMINSKI, M.A., GRADSTEIN, F.M. & BERGGREN, W.A. (1989). – Paleogene benthic foraminiferal stratigraphy and paleoecology at Site 647, Southern Labrador Sea. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 105, 705-730.
- KAMINSKI, M.A., GRADSTEIN, F.M., GOLL, R.M. & GRIEG, D. (1990). – Biostratigraphy and paleoecology of deep-water agglutinated foraminifera at ODP Site 643, Norwegian-Greenland Sea. – *In: Hemleben, C., Kaminski, M.A., Kuhn, W., & Scott, D.B. (eds), Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of Agglutinated Foraminifera*, NATO ASI Series, Kluwer Acad. Publ. 345-386.
- KAMINSKI, M.A., GRASSLE, J.F. & WHITLATCH, R.D. (1988). – Life History and recolonization among agglutinated foraminifera in the Panama Basin. – *In: Proceedings of the Second Workshop on Agglutinated Foraminifera, Vienna 1986. Abhandlungen der Geologischen Bundesanstalt*, 41, 229-244.
- KAMINSKI, M.A., KUHN, W. & RADLEY, J.D. (1996). – Palaeocene-Eocene deep water agglutinated foraminifera from the Numidian Flysch (Rif, Northern Morocco): their significance for the Palaeoceanography of the Gibraltar gateway. – *Journal of Micropalaeontology*, 15, 1-19.
- KAPTARENKO-CHERNOUSOVA, O.K., PLOTNIKOVA, L.F. & LIPNIK, E.S. (1979). – *Foraminifery Mela Ukrainy [Cretaceous Foraminifera of the Ukraine]*. – *Paleontologicheskii spravochnik*. Izdatelstvo "Naukova Dumka", Kiev, 256 pp.
- KARRER, F. (1866). – Über das auftreten von Foraminiferen in den älteren Schichten des Wiener Sandsteins. – *Sitzungsbereiche K. Akad. der Wissenschaften. Math.-Naturw. Classe Wien*, Bd. 52, (Jahrg 1865), Abt. 1, Heft 9, pp.
- KARRER, F. (1870). – Über ein neues Vorkommen von oberer Kreideformation in Leitersdorf bei Stockerau und deren Foraminiferenfauna. – *Jahrbuch der K.K. Geologischen Reichsanstalt*, 20, 157-184.
- KAVARY, E. & FRIZZEL, D.L. (1964). – Upper Cretaceous and lower Cenozoic foraminifera from west-central Iran. – *Missouri University School of Mines & Metallurgy Bulletin*, 102, 3-89.

- KENDER, S. (2003). – Stratigraphical significance of the foraminiferal assemblages from the Eocene Red Clays near Barwinek, described by Rudolf Noth in 1912 (Magura Unit, Outer Carpathians). – Unpublished M.Sc. report, Department of Geological Sciences, University College London, 83 pp.
- KHALILOV, D.M. (1959). – Novye vidy foraminifer Valanzhin-Goterivskikh otlozhenii severo-vostochnogo Azerbaidzhana [New species of foraminifera from the Valanginian-Hauterivian deposits of SE Azerbaidzan]. – *Izvestiya Akademii Nauk Azerbaidzhanskoi SSR, Seriya Geologo-Geograficheskikh Nauk*, 6, 25-36 + 2 pls.
- KING, C. (1983). – Cenozoic micropaleontological biostratigraphy of the North Sea. – *Institute of Geological Sciences Report*, 82/7, 1-40. London.
- KING, C. (1989). – Cenozoic of the North Sea. – *In: Jenkins, D.G. & Murray, J.W. (eds), Stratigraphical atlas of fossil foraminifera, second edition*, Ellis Horwood Ltd. 418-489.
- KING, C., BAILEY, H.D. & KING, A.D. (1989). – Cretaceous of the North Sea. – *In: Jenkins, D.G., and Murray, J.W. (eds), Stratigraphical atlas of fossil foraminifera, second edition*. Ellis Horwood, Ltd. 372-417.
- KINSEY, S. (1999). – Tertiary benthic foraminiferal biostratigraphy and palaeoecology of the Halten Terrace, Norway. – *GEOMAR Reports*, 77, 145 pp.
- KISSELMAN, E.N. (1964) – Otryad Textulariida. *In: Subbotina, N.N. (ed.), Foraminifery Melovykh I Paleogenovykh otlozhenii Zapadno-Sibirskoi Nizmennosti*. – Trudy VNIGRI, 234, 153-192.
- KOUTSOUKOS, E.A.M. (2000). – “Flysch-type” foraminiferal assemblages in the Cretaceous of northeastern Brazil. – *In: Hart, M.B., Kaminski, M.A. & Smart, C.W. (eds), Proceedings of the Fifth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 7, 243-260.
- KRAEVA, E.Y. & ZERNETSKY, BF. (1969). – *Foraminifery Paleogena Ukrainy [Paleogene Foraminifera of the Ukraine]*. – *Paleontologicheskii spravochnik*. Izdatelstvo "Naukova Dumka", Kiev, 197 pp + 76 pls.
- KRASHENNINIKOV, V.A. (1973). – Cretaceous benthonic foraminifera, Leg 20, Deep Sea Drilling Project. – *Initial Reports of the Deep Sea Drilling Project*, 20, 205-221.
- KRASHENNINIKOV, V.A. (1974). – Upper Cretaceous benthonic agglutinated foraminifera, Leg 27, Deep Sea Drilling Project. – *Initial Reports of the Deep Sea Drilling Project*, 27, 631-661.
- KRASHENNINIKOV, V.A. & PFLAUMANN, U. (1977). – Cretaceous agglutinated foraminifera of the Atlantic Ocean off West Africa (Leg 41, Deep Sea Drilling Project). – *Initial Reports of the Deep Sea Drilling Project*, 41, 565-580.
- KSIAŹKIEWICZ, M. (1975). – Bathymetry of the Carpathian flysch basin. – *Acta Geologica Polonica*, 25, 309-366.
- KUHNT, W. (1987a). – *Biostratigraphie und Palaoenvironment der externen Kreideserien des Westlichen Rif und Betikum - ein Ansatz zur Rekonstruktion der Kreide-Palaoeographie des Gibraltarbogens*. – Ph.D. Thesis, Eberhard- Karls-Universitat Tübingen. 271 pp.
- KUHNT, W. (1987b). – Upper Cretaceous foraminiferal assemblages of the external units of the Rif (Northern Morocco) - a paleobathymetric model of the Late Mesozoic North African continental margin. – *Géologie Méditerranéene*, 14 (2), 109-131.
- KUHNT, W. (1990). – Agglutinated foraminifera of western Mediterranean Upper Cretaceous pelagic limestones (Umbrian Apennines, Italy, and Betic Cordillera, Southern Spain). – *Micropaleontology*, 36, 297-330.
- KUHNT, W. & COLLINS, E.S. (1996). – Cretaceous to Paleogene benthic foraminifers from the Iberia Abyssal Plain. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 149, 203-216.
- KUHNT, W. & KAMINSKI, M.A. (1989). – Upper Cretaceous deep-water agglutinated benthic foraminiferal assemblages from the western Mediterranean and adjacent areas. – *In: Wiedmann, J. (ed.), Cretaceous of the Western Tethys. Proceedings of the 3rd International Cretaceous Symposium, Tübingen, 1987*. Schweitzerbart'sche Verlagsbuchhandlung, Stuttgart, 91-120.
- KUHNT, W. & KAMINSKI, M.A. (1990). – Paleoecology of Late Cretaceous to Paleocene deep-water agglutinated foraminifera from the North Atlantic and Western Tethys. – *In: Hemleben, C., Kaminski, M.A., Kuhnt, W. & Scott, D.B. (eds), Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of agglutinated foraminifera*. NATO ASI C, 327, 433-506. Kluwer Academic Press.
- KUHNT, W. & KAMINSKI, M.A. (1993). – Changes in the community structure of deep-water agglutinated foraminifers across the K/T Boundary in the Basque Basin (Northern Spain). – *Revista Española de Micropaleontología*, 25, 57-92.
- KUHNT, W. & KAMINSKI, M.A. (1996). – The response of benthic foraminifera to the K/T boundary event - a review. – *Bull. Centres Rech. Explor.-Production Elf Aquitaine Memoire*, 16, 433-442.

- KUHNT, W. & KAMINSKI, M.A. (1997). – Cenomanian to lower Eocene deep-water agglutinated foraminifera from the Zumaya section, northern Spain. – *Annales Societatis Geologorum Poloniae*, 67, 257-270.
- KUHNT, W. & URQUHART, E. (2001). – Tethyan flysch-type benthic foraminiferal assemblages in the North Atlantic: Cretaceous to Palaeogene deep water agglutinated foraminifera from the Iberia Abyssal Plain (ODP Leg 173). – *Revue de Micropaléontologie*, 44 (1), 27-59.
- KUHNT, W., COLLINS, E. & SCOTT, D.B. (2000). – Deep water agglutinated foraminiferal assemblages across the Gulf Stream: distribution patterns and taphonomy. – *In: Hart, M.B., Kaminski, M.A. & Smart, C.W. (eds), Proceedings of the Fifth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 7, 261-298.
- KUHNT, W., GEROCH, S., KAMINSKI, M.A., MOULLADE, M. & NEAGU, T. (1992). – Upper Cretaceous Abyssal Claystones in the North Atlantic and the Western Tethys - Current Status of Biostratigraphical Correlation using Agglutinated Foraminifera. – *Cretaceous Research*, 13, 467-478.
- KUHNT, W., HOLBOURN, A. & ZHAO, Q. (2002). – The early history of the South China Sea: Evolution of Oligocene–Miocene deep water environments. – *Revue de Micropaleontologie*, 45, 99-152.
- KUHNT, W., KAMINSKI, M.A. & MOULLADE, M. (1989). – Deep-water agglutinated benthic foraminiferal assemblages of the upper Cretaceous North Atlantic and its marginal seas. – *Geologische Rundschau*, 78 (3), 1121-1140.
- KUHNT, W., MORLOTTI, E., WINKLER, W. & KAMINSKI, M.A. (1989). – *Late Cretaceous to early Paleogene Alpine sequences and their agglutinated foraminifera*. – NATO Advanced Studies Workshop Series. Excursion Guidebook, 156 pp.
- KUHNT, W., MOULLADE, M. & KAMINSKI, M.A. (1998). – Upper Cretaceous, K/T boundary, and Paleocene agglutinated foraminifera from Hole 959D (Côte d'Ivoire – Ghana transform Margin). – *Proceedings of the Ocean Drilling Program, Scientific Results*, 159, 389-411.
- LABUDE, C. (1984). – Sedimentologie, Subsidenz und Verbreitung sandschaliger und planktonischer Foraminiferen im Maastricht des Zentralen Apennin (Italien). – *Tübinger Mikropaläontologische Mitteilungen*, 2, 95 pp.
- LALICKER, C.G. (1935). – New Tertiary Textulariidae. – *Contributions from the Cushman Laboratory for Foraminiferal Research*, 11, 1-13.
- LANGER, M.R., LIPPS, J.H. & MORENO, G. (1995). – Predation on foraminifera by the dentaliid deep-sea scaphopod *Fissidentalium megathyris*. – *Deep-Sea Research Part I-Oceanographic Research Papers*, 42 (6), 849-857.
- LAURSEN, G.V. & ANDERSEN, S.B. (1997). – A Late Palaeocene - Early Eocene benthic foraminiferal record from Bovlstrup, Denmark, showing a remarkable agglutinated fauna – *Journal of Micropalaeontology*, 16, 19-29
- LEROY, L.W. (1964). – Smaller foraminifera from the Late Tertiary of southern Okinawa. *U.S. Geological Survey Professional Paper*, 454-F, 58 pp.
- LEŠKO B. & SAMUEL O. (1968). – *Geologia východoslovenského flyšu*. – Vydavateľstvo Slov. Akad. Vied, Bratislava. 245 pp.
- LEVIN, L., BLAIR, N., DEMASTER, D., PLAIA, G., FORNES, W., MARTIN, C. & THOMAS, C. (1997). – Rapid subduction of organic matter by maldanid polychaetes on the North Carolina slope. – *Journal of Marine Research*, 55, 595-611.
- LIEBUS A. & SCHUBERT J.R. (1903). – Die Foraminiferen der Karpathischen Inoceramenschichten von Gbellan in Ungarn (Puchower Mergel). – *Jahrbuch der. K.-kön. geologischen Reichsanstalt (Wien)*, 52, 2, 285-310.
- LINDENBERG, H.G. (1967). – Die Arten von *Haplophragmium* und *Triplasia*. – *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, 514, 1- 74. Frankfurt am Main.
- LINDENBERG, H.G. & AURAS, A. (1984). – Distribution of arenaceous foraminifera in depth profiles of the Southern Ocean. – *Paleogeography Paleoclimatology Paleoecology*, 48, 61-106.
- LINKE, P. (1992). – Metabolic adaptations of deep sea benthic foraminifera to seasonally varying food input. – *Marine Ecology-Progress Series*, 81 (1), 51-63.
- LINKE, P. & LUTZE, G.F. (1993). – Microhabitat preferences of benthic foraminifera - a static concept or a dynamic adaptation to optimize food acquisition? – *Marine Micropaleontology*, 20, 215-234.
- LIPPS, J.H. & VALENTINE, J.W. (1970). – The role of foraminifera in the trophic structure of marine communities. – *Lethaia*, 3, 279-286.
- LISZKA, S. & LISZKOWA, J. (1981). – Revision of J. Grzybowski's paper (1896) "Foraminifera of the red clays from Wadowice". – *Rocznik Polskiego Towarzystwa Geologicznego*, 51, 153-208.

- LOEBLICH, A.F. (1946). – Foraminifera from the type Pepper Shale of Texas. – *Journal of Paleontology*, 20, 130-139.
- LOEBLICH, A.F. & TAPPAN, H. (1964). – Sarcodina chiefly "Thecamoebians" and Foraminiferida. – In: Moore, R.C. (ed.), *Treatise on invertebrate paleontology, part C, Protista*. University of Kansas Press. 900 pp.
- LOEBLICH, A.R. & TAPPAN, H. (1984). – Some new proteinaceous and agglutinated genera of Foraminiferida. – *Journal of Paleontology*, 58, 1158-1163.
- LOEBLICH, A.R. & TAPPAN, H. (1987). – *Foraminiferal genera and their classification*. – Van Nostrand Reinhold Co., 970 pp.
- LOEBLICH, A.R. & TAPPAN, H. (1989). – Implications of wall composition and structure in agglutinated foraminifers. – *Journal of Paleontology*, 63, 769-777.
- LOWRIE, W., ALVAREZ, A., NAPOLEONE, G., PERCH-NIELSEN, K., PREMOLI SILVA, I. & TOUMARKINE, M. (1982). – Paleogene magnetic stratigraphy in the Umbrian pelagic carbonate rocks: The Contessa sections, Gubbio. – *Geological Society of America Bulletin*, 93, 414-432.
- LUDBROOK, N.H. (1977). – Early Tertiary *Cyclammmina* and *Haplophragmoides* (Foraminiferida: Lituolidae) in southern Australia. – *Transactions of the Royal Society of South Australia*, 101, 165-198.
- LUKINA, T.G. (1980). – Globokovodnye foraminifery tsentralnoy chasti Tikhogo Okeana [Deep water foraminifera from the central part of the Pacific Ocean]. – *Issledovaniya Fauny Morey*, 24, Izdatel'stvo Nauka, Leningrad, 203 pp.
- LUTERBACHER, H.P. & PREMOLI SILVA, I., 1964. – Biostratigrafia del limite Cretaceo-Terziario nell'Appennino centrale. – *Rivista Italiana de Paleontologia e Stratigrafia*, 70, 67-128.
- MacFADYEN, W.A. (1933). – Fossil foraminifera from the Burdwood Bank and their geological significance. – *Discovery Reports*, 7, 3-16.
- MacFADYEN, W.A. (1933). – A note on the foraminiferal genus *Bolivinopsis* Yakovlev. – *Journal of the Royal Microscopical Society*, 53, 139-141.
- MACKENSEN, A., SCHMIEDL, G., HARLOFF, J. & GIESE, M. (1995). – Deep-sea foraminifera in the south Atlantic Ocean: Ecology and assemblage generation. – *Micropaleontology*, 41, 342-358
- MAJZON, L. (1943). – Adatok egyes Kárpátaljai Flisrétegekhez, tekintettel a Globotruncanákra [Beitrage zur Kenntniss einiger Flysch-Schichten des Karpaten-Vorlandes mit Rucksicht auf die Globotruncanen]. – *A magyar Királyi Földtani Intézet, Évkönyve*, 37, 1-170.
- MALATA, E. & OSZCZYPKO, N. (1990). – Deep water agglutinated foraminiferal assemblages from Upper Cretaceous red shales of the Magura Nappe / Polish Outer Carpathians. – In: Hemleben, C., Kaminski, M.A., Kuhnt, W. & Scott, D.B. (eds), *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of agglutinated foraminifera*. NATO ASI C, 327, 507-524. Kluwer Academic Press.
- MALLORY, V.S. (1959). – *Lower Tertiary biostratigraphy of the California Coastal Ranges*. – American Association of Petroleum Geologists, 416 pp.
- MARKS, P. (1951). – A revision of the smaller foraminifera from the Miocene of the Vienna Basin. – *Contributions from the Cushman Foundation for Foraminiferal Research*, 2, 33-73.
- MARTIN, L. (1964). – Upper Cretaceous and Lower Tertiary foraminifera from Fresno County, California. – *Jahrbuch der Geologischen Bundesanstalt (Austria), Sonderband*, 9, 1-128.
- MARSSON, T. (1878). – Die Foraminiferen der weissen Schreibkreide der Insel Rugen. – *Mitteilungen des Naturwissenschaftlichen Verein für Neu-Vorpommern und Rugen in Greifswald*, 10, 115-196.
- MASLAKOVA, N.I. (1955). – Stratigrafiya i fauna melkikh foraminifer paleogenovykh otlozheniy Vostochnykh Karpat. – *Materialy po Biostratigrafi zapadnykh oblastii Ukrainskoi SSR*, 5-132. Moskva.
- MASLUN, N.V. (1987). – Agglutinirovannyye foraminifery pogranichnykh Eocene-Oligotsenovykh otlozheniy Karpat [Agglutinated foraminifera from the Eocene/Oligocene boundary beds of the Carpathians]. – *Paleontologicheskij Sbornik, L'viv*, 27, 66-70. Lvov.
- MATSUNAGA, T. (1955). – *Spirosigmoinella*, a new foraminiferal genus from the Miocene of Japan. – *Transactions and Proceedings of the Palaeontological Society of Japan, n.s.*, Tokyo, 18, 49-50.
- McCABE, C.M., KAMINSKI, M.A. & FINCH, E. (1993). – A biostratigraphic revision of the Eocene and Oligocene type localities of Trinidad described by Cushman and Stainforth (1945) and Cushman & Renz (1948). – *Journal of Micropalaeontology*, 12, 195-200.
- McDOUGALL, K. (1980). – Paleocological evaluation of late Eocene biostratigraphic zonations of the Pacific coast of North America. – *Society of Economic Paleontologists and Mineralogists Paleontological Monographs*, 2 (Journal of Paleontology vol. 54 supplement) 75 pp.
- McNEIL, D.H. (1989). – Foraminiferal zonation and biofacies analysis of the Cenozoic strata in the Beaufort-MacKenzie Basin of Arctic Canada. – *Current Research, Part G, Geological Survey of Canada Paper*, 89-1G, 203-233.

- McNEIL, D.H. (1990). – Tertiary marine events of the Beaufort-Mackenzie Basin and correlation of Oligocene to Pliocene marine outcrops in Arctic North America. – *The Arctic*, 43 (4), 301-313.
- McNEIL, D.H. (1996). – Distribution of Cenozoic agglutinated benthic foraminifera in the Beaufort-Mackenzie Basin. – In: Dixon, J. (ed.) Geological Atlas of the Beaufort-Mackenzie Area. *Geological Survey of Canada Miscellaneous Report*, 59, fig. 70.
- McNEIL, D.H. (1997). – New foraminifera from the Upper Cretaceous and Cenozoic of the Beaufort-Mackenzie Basin of Arctic Canada. – *Cushman Foundation for Foraminiferal Research Special Publication*, 35, 95 pp.
- McNIEL, D.H. & CALDWELL, W.G.E. (1981). – Cretaceous rocks and their foraminifera in the Manitoba Escarpment. – *Geological Association of Canada Special Paper*, 21, 439 pp.
- McNULTY, C.L. (1979). – Smaller Cretaceous foraminifers of Leg 43, Deep Sea Drilling Project. – *Initial Reports of the Deep Sea Drilling Project*, 43, 487-505.
- MELLO, J.F. (1971). – Foraminifera from the Pierre Shale (Upper Cretaceous) at Red Bird, Wyoming. – *U.S. Geological Survey Professional Paper*, 393-C, 54 pp.
- MILLER, K.G. & KATZ, M. (1987). – Oligocene to Miocene benthic foraminiferal and abyssal circulation changes in the North Atlantic. – *Micropaleontology*, 33, 97-149.
- MILLER, K.G., GRADSTEIN, F.M. & BERGGREN, W.A. (1982). – Late Cretaceous to early Tertiary agglutinated benthic foraminifera in the Labrador Sea. – *Micropaleontology*, 28, 1-30.
- MILLETT, F.W. (1899a). – Report on the Recent Foraminifera of the Malay Archipelago collected by Mr. A. Durrand, F.M.R.S., part IV. – *Journal of the Royal Microscopical Society*, 1988, 249-255.
- MILLETT, F.W. (1899b). – Report on the Recent Foraminifera of the Malay Archipelago collected by Mr. A. Durrand, F.M.R.S., part V. – *Journal of the Royal Microscopical Society*, 1988, 357-365.
- MILNER, G.J. (1997). – *Paleogene foraminiferal studies in the Western Pacific. Part A. Paleocene foraminiferal biostratigraphy of the Papuan Basin and adjoining areas, Papua New Guinea; Part B. Middle Eocene to early Oligocene foraminifers from the Izu-Bonin fore-arc, Hole 786A.* – Unpublished Ph.D. dissertation, University of Western Australia, 302 pp.
- MJATLIUK, E.V. (1950). – Stratigrafiya flishevykh osadkov Severnykh Karpat w svete dannykh fauny foraminifer [Stratigraphy of the flysch deposits of the northern Carpathians in the light of foraminiferal data]. – *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Instituta VNIGRI*, 51, 225-287. Moskva.
- MJATLIUK, E.V. (1966). – K voprosu o foraminiferakh s kremnevnyim skeletom [On the question of foraminifera with a siliceous skeleton]. – *Voprosy Mikropaleontologii*, 10, 255-269.
- MJATLIUK, E.V. (1970). – Foraminifery flishevykh otlozhenii vostochnykh Karpat (Mel-Paleogen) [Foraminifera of the flysch deposits of the eastern Carpathians]. – *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Instituta VNIGRI*, 282, 1-225. Leningrad.
- MONTANARI, A. & KOEBERL, C. (2000). – Impact Stratigraphy, The Italian Record. – *Lecture Notes in Earth Sciences*, 93, 349 pp. Springer-Verlag, Berlin.
- MONTANARO GALLITELLI, E. (1955). – Foraminiferi cretacei delle marne a fucoidi di Serramazzoni (Appennino modenese). – *Accad. Sci. Lett. Arti Modena, Atti Mem., Modena Italy, ser. 5*, vol. 143.
- MOORKINS, T.L. (1976). – Palökologische Bedeutung einiger Verschellschaftungen von sandschaligen Foraminiferen aus dem NW europäischen Alttertiär und ihre Beziehung zu Muttergesteinen. – *Erdöl und Kohle, Erdgas, Petrochemie, Compendium*, 75/76, 77-95.
- MORENO-VÁSQUEZ, J. (1995). – Neogene biofacies in eastern Venezuela and their calibration with seismic data. – *Marine Micropaleontology*, 26 (1-4), 287-302.
- MORGIEL, J. & OLSZEWSKA, B. (1981). – Biostratigraphy of the Polish External Carpathians based on agglutinated foraminifera. – *Micropaleontology*, 27, 1- 30.
- MORGIEL, J. & OLSZEWSKA, B. (1982). – Uniformity of the Tethyan faunas from Cretaceous and Paleogene as shown by foraminifera from Morocco and Polish Flysch Carpathians. – *Cahiers de Micropaleontologie*, 3, 45-53.
- MORGIEL, J. & SZYMAKOWSKA, F. (1978). – Stratygrafia paleocenu i eocenu jednostki skolskiej [Stratigraphy of the Paleocene and Eocene of the Skole Unit]. – *Biuletyn Instytutu Geologicznego*, 310, 39-71 + 15 pls.
- MORLOTTI, E. & KUHN, W. (1992). – Agglutinated deep-water foraminifera of the Eocene Monte Piano Formation (Northern Apennines, Italy). – *Journal of Foraminiferal Research*, 22, 214-228.
- MORRIS, R.W. (1971). – Upper Cretaceous Foraminifera from the upper Mancos Formation, northwestern Colorado. – *Micropaleontology*, 17, 257-296.

- MOULLADE, M., KUHNT, W. & THUROW, J. (1988). – Agglutinated benthic foraminifers from Upper Cretaceous variegated clays of the North Atlantic Ocean. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 103, 349-377.
- MUDGE, D.C. & BUJAK, J.P. (1996). – An integrated stratigraphy for the Paleocene and Eocene of the North Sea. – In: Knox, R.W.O'B., Corfield, R.M. & Dunay, R.E. (eds), *Correlation of the Early Paleogene in Northwest Europe*. Geological Society Special Publication, 101, 91-113.
- MURRAY, J.W. & ALVE, E. (1994). – High diversity agglutinated foraminiferal assemblages from the NE Atlantic dissolution experiments. – *Cushman Foundation Special Publication*, 32, 33-51.
- MÜLLER-MERZ, E. & OBERHÄNSLI, H. (1991). – Eocene bathyal and abyssal benthic foraminifera from a South Atlantic transect at 20–30° S. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 83, 117-171.
- NAGY, J. (1992). – Environmental significance of foraminiferal morphogroups in Jurassic North Sea deltas. – *Palaeogeography Palaeoclimatology Palaeoecology*, 95 (1-2), 111-134.
- NAGY, J., GRADSTEIN, F.M., KAMINSKI, M.A. & HOLBOURN, A.E.L. (1995). – Late Jurassic to Early Cretaceous foraminifera of Thakkhola, Nepal: Palaeoenvironments and description of new taxa. – In: Kaminski, M.A., Geroch, S. & Gasinski, M.A. (eds), *Proceedings of the Fourth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 3, 181-209.
- NAGY, J., KAMINSKI, M.A., JOHNSEN, K. & MITLEHNER, A.G. (1997). – Foraminiferal, palynomorph, and diatom biostratigraphy and palaeoenvironments of the Torsk Formation: A reference section for the Paleocene-Eocene transition in the western Barents Sea. – In: Hass, H.C. & Kaminski, M.A. *Contributions to the Micropaleontology & Paleoceanography of the Northern North Atlantic*. Grzybowski Foundation Special Publication, 5, 15-38.
- NAGY, J., KAMINSKI, M.A., KUHNT, W. & BREMER, M.A. (2000). – Agglutinated foraminifera from neritic to bathyal facies in the Palaeogene of Spitsbergen and the Barents Sea. – In: Hart, MB; Kaminski, MA; Smart, CW (eds), *Proceedings of the Fifth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 7, 333-361.
- NAGY, J., KAMINSKI, M.A., GRADSTEIN, F.M. & JOHNSON, K. (2004). – Quantitative foraminiferal and palynomorph biostratigraphy of the Paleogene in the southwestern Barents Sea. – In: Bubík, M. & Kaminski, M.A., (eds), *Proceedings of the Sixth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 8, 359-379.
- NAUSS, A.W. (1947). – Cretaceous microfossils of the Vermilion area, Alberta. – *Journal of Paleontology*, 21, 329-343.
- NEAGU, T. (1962). – Studiul foraminiferelor aglutinante din argilele Cretacic superioare de pe Valea Sadovei (Cîmpulung-Moldovenesc) si bazinul superior al vaii Buzauliu. – *Studii Cercetari Geologie, Academia Republicii Populare Romine, Sectia de Geologie si Geografie si Institutul de Geologie si Geografie*, 7, 45-81.
- NEAGU, T. (1964). – Large size agglutinated foraminifera from the Carpathian of Rumania. – *Rocznik Polskiego Towarzystwa Geologicznego*, 34, 579-588.
- NEAGU, T. (1970). – Micropaleontological and stratigraphical study of the upper Cretaceous deposits between the upper valleys of the Buzau and Riul Negru Rivers (Eastern Carpathians). – *Memorii, Institutul Geologic*, 12, 7-109. Bucarest.
- NEAGU, T. (1972). – The Eo-Cretaceous foraminiferal fauna from the area between the Lalomitza and Prahova valleys (Eastern Carpathians). – *Revista Española de Micropaleontologia*, 4 (2), 181-224.
- NEAGU, T. (1975). – Monographie de la faune des Foraminifères éocétacés du couloir de Dimboviciora, de Codlea et des Monts (Persani), (Couches de Carhaga). – *Memorii, Institutul Geologic*, 25, 1-141.
- NEAGU, T. (1990). – *Gerochammina* n.gen. and related genera from the Upper Cretaceous flysch-type benthic foraminiferal fauna, Eastern Carpathians – Romania. – In: Hemleben, C., Kaminski, M.A., Kuhnt, W., & Scott, D.B. (eds), *Paleoecology, Biostratigraphy, Paleoceanography and Taxonomy of agglutinated foraminifera*. NATO ASI C, 327, 245-265. Kluwer Academic Press.
- NEAGU, T & PLATON, E. (1994). – Genera *Haplophragmoides* Cushman, 1910; *Recurvoides* Earland, 1934; *Thalmannammina* Pokorny, 1951; *Plectorecurvoides* Noth, 1952; and *Pokornyammina* n.gen. from Upper Cretaceous flysch facies, Eastern Carpathians, Romania. – *Revista Española de Micropaleontologia*, 26 (1), 5-30.
- NIKITINA, Yu.P. & SVEMBERGER, Yu.N. (1963). – Paleotsen Azovo-Kubanskoy vpadiny, Vala Karpinskogo i Yuzhnogo sklona Vorozhskoy Anteklizy. – In: Burshmara, M.S. (ed.), *Materialy po Geologii i Neftogazonosnosti Yuga SSSR*, Trudy VNIGRI, 38, 105-147. Gosgeoltechizdat, Moscow.
- NISHIDA, T., MATSUMOTO, T. & INOUE, Y. (1998). – Foraminifera from the Cretaceous System in the Shumarinai Valley of Hokkaido. – *Journal of the Faculty of Culture and Education, Saga University*, 3 (1), 301-331.

- NORTH, B.R. & CALDWELL, W.G.E. (1964). – Foraminifera from the Cretaceous Lea Park Formation in south-central Saskatchewan. – *Saskatchewan Research Council, Geology Division Report*, 5, 43 pp.
- NØRVANG, A. (1941). – Notes on some foraminifera from off Bergen. – *Bergens Museum Årbok, Naturvitens rekke*, 11, 1-19.
- NOÉ, S.U. (1993). – Zyklusstratigraphie und Foraminiferen-Palökologie in pelagoschen Sedimentationzyklen des Apt/Alb im Umbrisch-Marchischen Apennin (Mittelitalien). – *Zitteliana*, 20, 179-211.
- NOTH, R. (1912). – Die Foraminiferenfauna der roten Tone von Barwinek und Karmarnók. – *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients*, 25, 1-24 + 1 pl.
- NOTH, R. (1951). – Foraminiferen aus Unter- und Oberkreide des österreichischen Anteils an Flysch, Helvetikum und Vorlandvorkommen. – *Jahrbuch der Geologischen Bundesanstalt, Sonderband*, 3, 1-91.
- NUTTALL, W.L.F. (1930). – Eocene foraminifera from Mexico. – *Journal of Paleontology*, 4 (3), 271-293.
- OLSZEWSKA, B. (1983). – A contribution to the knowledge of planktonic foraminifers of the Globigerina submenilite marls in the Polish Outer Carpathians [in Polish]. – *Kwartalnik Geologiczny*, 27, 547-570.
- OLSZEWSKA, B. (1984). – Interpretacja paleoekologiczna otwornic kredy i paleogenu polskich Karpat zewnętrznych [Paleoecological interpretation of Cretaceous and Paleogene foraminifera of the Polish external Carpathians]. – *Biuletyn Instytutu Geologicznego*, 346, 7-62.
- OLSZEWSKA, B. (1997). – Foraminiferal biostratigraphy of the Polish Outer Carpathians: a record of basin geohistory. – *Annales Societatis Geologorum Poloniae*, 67, 325-337.
- OLSZEWSKA, B. & SMAGOWICZ, M. (1977). – Porównanie podziałów biostratygraficznych górnej kredy i paleogenu jednostki dukielskiej na podstawie otwornic planktonicznych i nannoplantonu wapiennego [A comparison of Upper Cretaceous and Paleogene biostratigraphic schemes based on planktonic foraminifera and nannofossils]. – *Przegląd Geologiczny*, 7, 359-363.
- ORTIZ, N. (1995). – Differential patterns of benthic foraminiferal extinctions near the Paleocene/Eocene boundary in the North Atlantic and the western Tethys. – *Marine Micropalontology*, 26, 341-359.
- ORUE EXTEBARRIA X., APELLANIZ E., BACETA J.J., COCCIONI R., DI LEO R., DINARES-TURELL J., GALEOTTI S., MONECHI S., NUÑEZ -BETELU K., PARES J.M., PAYROS A., PUJALTE V., SAMSO J.P., SERRA-KIEL J., SCHMITZ B. & TOSQUELLA, J., (1996). – Physical and biostratigraphic analysis of two prospective Paleocene-Eocene Boundary Stratotypes in the intermediate-deep water Basque Basin, western Pyrenees: The Trabakua Pass and Ermua sections. – *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 201, 179-242.
- OSTERMAN, L.E. & QUALE, G. (1989). – Benthic foraminifers from the Vøring Plateau (ODP Leg 104). – *Proceedings of the Ocean Drilling Program, Scientific Results*, 104, 745-768.
- OSTERMAN, L. & SPIGLER, D. (1996). – Agglutinated benthic foraminiferal biostratigraphy of Ocean Drilling Program (ODP) Sites 909 and 913, northern North Atlantic. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 151, 169-185.
- PAPP, A; SCHMID, ME. (1985). – The fossil Foraminifera of the Tertiary Basin of Vienna: Revision of the monograph by Alcide d'Orbigny (1846). – *Abhandlungen der geologischen Bundesanstalt*, 37, 1-311.
- PARKER, F.L. (1952). – Foraminiferal species off Portsmouth, New Hampshire. – *Bulletin of the Museum of Comparative Zoology*, 106 (9), 391-423 + 6 pls.
- PARKER, F.L. (1964). – Foraminifera from the experimental Mohole drilling near Guadalupe Island, Mexico. – *Journal of Paleontology*, 38, 617-636.
- PARR, W.J. (1950). – Part 6, Foraminifera. – *B.A.N.Z. Antarctic Research Expedition 1929-1931 Reports, Series B (Zoology and Botany)*, vol. 5, part 6, 237-392.
- PERYT, D, LAHODYNSKY, R & DURAKIEWICZ, T. (1997). – Deep-water agglutinated foraminiferal changes and stable isotope profiles across the Cretaceous-Paleogene boundary in the Rotwandgraben section, Eastern Alps (Austria). – *Palaeogeography Palaeoclimatology Palaeoecology*, 123 (1-4), 287-307.
- PETRACCA, A.N. (1972) – Tertiary microfauna, MacKenzie Delta area, Arctic Canada. – *Micropaleontology*, 18, 355-368.
- PETERSON, R.H., GAUGER, D.J. & LANKFORD, R.R. (1953). – Microfossils of the Upper Cretaceous of northeastern Utah and southwestern Wyoming. – *Utah Geology and Mining Survey Bulletin*, 47, 158 pp.
- PFLAUMANN, U. (1964). – *Geologisch-mikropalaontologische Untersuchungen in der Flysch-Oberkreide zwischen Wertach und Chiemsee in Bayern*. – Inaugural dissertation. Ludwig Maximilian Universität. München, 180 pp.
- PFLAUMANN, U. (1967). – Zur Ökologie des bayerischen Flysches auf Grund der Mikrofossilführung. – *Geologisches Rundschau*, 56, 200-227.

- PFLAUMANN, U. (1968). – II. Flysch-Zone. – *Erläuterungen zur Geologischen Karte von Bayern 1:25 000 Blatt Nr. 8237 Miesbach*. 111- 415.
- PFLAUMANN, U. (1973). – ???(Cenoman am Tiefbajuvarikum-Nordrand.) – *Erläuterungen zu Geologische Karte von Bayern 1:25 000 Blatt Nr. 8238 Neubeuern*. 114-150 pp.
- PFLUM, C.E. (1966). – The distribution of foraminifera in the Eastern Ross Sea, Amundsen Sea, and Bellingshausen Sea, Antarctica. – *Bulletins of American Paleontology*, 50 (226), 151-209.
- PFLUM, C.E. & FRERICHS, W.E. (1976). – Gulf of Mexico deep water foraminifers. – *Cushman Foundation for Foraminiferal Research Special Publication*, 14, 1-125.
- PHLEGER, F.B (1951). – Ecology of Foraminifera, northwest Gulf of Mexico. Part 1, Foraminifera distribution. – *Geological Society of America Memoir*, 46, 88 pp.
- PHLEGER, F.B (1952). – Foraminifera ecology off Portsmouth, New Hampshire. – *Bulletin of the Museum of Comparative Zoology*, 106 (8), 315-390.
- PHLEGER, F.B & PARKER, F.L. (1951). – Ecology of Foraminifera, northwest Gulf of Mexico, Part 2, Foraminiferal species. – *Geological Society of America Memoir*, 46, 64 pp.
- PLATON, E (1997). – Coiling modes in the family Plectrorecurvoididae (Foraminiferida). – *Annales Societatis Geologorum Poloniae*, 63 (2-3), 339-342.
- POAG, C.W. (1981). – *Ecologic Atlas of benthic foraminifera of the Gulf of Mexico*. – Stroudsburg, PA (Hutchinson Ross Publishing Co.), 174 pp.
- PODOBINA, V. (1966). – *Foraminifery vyerkhnego mela Zapadno-sibirskoy Nizmennosti*. – Izdatel'stvo Nauka, Moscow, 148 pp + 19 pls.
- PODOBINA, V. (1975). – *Foraminifera of the Upper Cretaceous and Paleocene of the western Siberian lowlands, and their significance for stratigraphy*. – Izdatel'stvo Tomskogo Universiteta, Tomsk. 220 pp + 40 pls. [in Russian].
- POKORNÝ, V. (1949). – On the microfauna of the Eocene Green Clay of Nikolčice (Moravia, Czechoslovakia). – *Bulletin International de l'Académie tchèque des Sciences* 1949, 15, 1-17.
- POKORNÝ, V. (1951). – *Thalmanammia* n.g. (Foraminifera) from the Carpathian flysch. – *Sborník Ústředního Ústavu Geologického*, 18, 469-480. Praha.
- POKORNÝ, V. (1953). – Mikrobiostratigrafická posice herspických sterku v eocénu zdánické serie (s popisem foraminifer sousedních jílů. – *Rozpravy II. Trídy České Akademie*, 61, (28), 1-36.
- POKORNÝ, V. (1960). – Microstratigraphie et biofaciès du flysch Carpatique de la Moravie Méridionale (Tchécoslovaquie). – *Revue de l'Institut Français du Pétrole et Annales des Combustibles Liquides*, 15, 1099-1141.
- POPESCU, G. (1972). – Subfamilia Cyclammininae (Foraminiferida) în Oligo-Miocenul din Transilvania de Nord. – *Studii si Ceretari de Geologie Geofizica Geografie, seria Geologie*, 17, 299-321.
- POPESCU, G. & IVA, M. (1971). – Contribution á la connaissance de la microfauna Oligocène des couches de Valea Lapusului. – *Memorii, Institutul Geologic*, 14, 35-51.
- PREECE, R.C. (1999). – *The physiological response of equatorial Neogene bathyal benthic foraminifera to low oxygen conditions*. – Unpublished Ph.D. Thesis, University of London, 375 pp + 84 pls.
- PREECE, R.C., KAMINSKI, M.A. & DIGNES, T.W. (1999). – Miocene benthonic foraminiferal morphogroups in an oxygen minimum zone, offshore Cabinda. – In: Cameron, N.R. et al. (eds), *The Oil and Gas Habitats of the South Atlantic. Geological Society (London), Special Publication*, 153, 267-282.
- PROTO-DECIMA, F. & BOLLI, H.M. (1978). – Southeast Atlantic DSDP Leg 40, benthic foraminifera. – *Initial Reports of the Deep Sea Drilling Project*, 40, 783-809.
- REISER, H. (1987). – Die Foraminiferen der bayerischen Oligozän-Molasse Systematic, Stratigraphie und Paläobathymetrie. – *Zitteliana*, 16, 3-131 + 19 pls.
- REUSS, A.E. (1845). – *Die Versteinerungen der böhmischen Kreideformation*. – Abtheilung 1. E. Schweizerbart'sohne Verlagsbuchhandlung, Stuttgart. 58 pp.
- REUSS, A.E. (1846). – *Die Versteinerungen der böhmischen Kreideformation (Nachträge und Ergänzungen zur ersten Abtheilung), Abtheilung 2*. – E. Schweizerbart'sohne Verlagsbuchhandlung, Stuttgart. 148 pp.
- REUSS, A.E. (1851). – Über die fossilen Foraminiferen und Entomostraceen der Septarianthone der Umgegend von Berlin. – *Deutsche Geologische Gesellschaft, Zeitschrift, Berlin*, 3, 49-91.
- REUSS, A.E. (1860). – Die Foraminiferen der westphalischen Kreideformation. – *Sitzungsberichte der Kaiserliche Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse*, 40, 147-238.

- REUSS, A.E. (1862). – Entwurf einer systematischen Zusammenstellung der Foraminiferen. *Sitzungsberichte der Kaiserliche Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse*, 44, 355-396.
- RHUMBLER, L. (1895). – Entwurf eines natürlichen Systems der Thalamophoren. – *Nachrichten der K. Gesellschaft der Wissenschaften zu Göttingen, mathematisch-physikalische Klasse*, 1, 4-98.
- RHUMBLER, L. (1903). Systematische Zusammenstellung der recenten Reticulosa (Nuda + Foraminifera). – *Archiv für Protistenkunde*, 1903, 181-294.
- RHUMBLER, L. (1905). – Mitteilungen über Foraminiferen (mit Demonstrationen). – *Verhandlungen der Deutschen Zoologischen Gesellschaft*, 15, 97-106.
- RHUMBLER, L. (1913) – Die Foraminiferen (Thalamophoren) der Plankton-Expedition, zwierter Teil, Systematik: Arrhabdammidia, Arammodiscidia und Arnodosammidia. – *Ergebnisse der Plankton-Expedition der Humboldt-Stiftung, Kiel und Leipzig, Band 3*, 332-476.
- RIEGRAF, W. & LUTERBACHER, H.-P. (1989). – Benthic foraminifera from the Lower Cretaceous (Deep Sea Drilling Project Leg 1-79). – *Geologische Rundschau*, 78, 1063-1120.
- ROBERTSON, D. (1891). – *Trochammina bradyi* n.n. – *Ann. Mag. Nat. Hist. London, ser. 6, 7*, 388.
- RÖGL, F. (1976). – Late Cretaceous to Pleistocene foraminifera from the Southeast Pacific Basin, DSDP Leg 35. – *Initial Reports of the Deep Sea Drilling Project*, 35, 539-555.
- RÖGL, F. (1995). – A Late Cretaceous flysch-type agglutinated foraminiferal fauna from the *Trochamminoides proteus* type locality (Wien - Hütteldorf, Austria) - In: Kaminski, M.A., Geroch, S. & Gasinski, M.A. (eds), *Proceedings of the Fourth International Workshop on Agglutinated Foraminifera*. Grzybowski Foundation Special Publication, 3, 249-263.
- RÖGL, F., FUCHS, R., SCHNABEL, W., SEIFERT, P. & WAGNER, L. (1986). – *Excursion Guide to the 2nd IWAF Excursion in Austria 1986*. Second International Workshop on Agglutinated Foraminifera, Abstracts, Program and Excursion Guide. pp. 60-92. H. Peter Press. Strasshof, Austria.
- RZEHAK, A. (1887). – Die Foraminiferenfauna des grünen Oligocenthones von Nikolschitz. – *Verhandlungen der Geologischen Reichsanstalt in Wien*, 1888. pp. 87-88.
- RZEHAK, A. (1895). – Über einige merkwürdige Foraminiferen aus dem Österreichischen Tertiär. – *Annalen des K.K. Naturhistorisches Hofmuseum*, 10, 213-230. Wien.
- SAID, R. & KENAWY, A. (1956). – Upper Cretaceous and lower Tertiary foraminifera from northern Sanai, Egypt. – *Micropaleontology*, 2, 105-173.
- SAIDOVA, Kh. M. (1961). – *Ekologia foraminifer i paleogeografiya dal'nevostochnykh chasti SSSR i severno-zapadnoy chasti Tikhogo Okeana* [Foraminiferal ecology and paleogeography of the far-eastern part of the USSR and the northwest part of the Pacific Ocean] - Izdat. Akad. Nauk SSSR, Moskva, 232 pp.
- SAIDOVA, Kh. M. (1975). – *Bentosnye Foraminifery Tikhogo Okeana* [Benthonic foraminifera of the Pacific Ocean]. – Institut Okeanologii P.P. Shirshova, 3. Akademiya Nauk SSSR, Moscow.
- SAMUEL, O. (1977). – Agglutinated foraminifera from Paleogene flysch formations in West Carpathians of Slovakia. – *Zapadne Karpaty, serie paleontologie*, 2-3, 7-70.
- SANDULESCU, J. (1972). – Étude micropaléontologique et stratigraphique du flysch du Crétacé supérieur-Paléocène de la région de Bretcu-Comandau (Secteur intern meridional de la nappe de Tarcau-Carpates Orientales). – *Memorii, Institutul Geologic*, 17, 1-52. Bucarest.
- SANDULESCU, J. & BRATU, E. (1981). – Nouvelles données pour la corrélation des Formations Paléogènes des unités allochtones de la Zone du Flysch Transcarpathique (Maramures). – *Dari de Seama Institutului de Geologie si Geofizica*, 68 (4), 199-135.
- SANSORES & FLORES-COVARRUBIAS, (1972). – Descripción de seis nuevas especies de foraminiferos bentónicos de la Cuenca Salina del Istmo de Tehuantepec, Mexico. – *Rev. del Inst. Mex. del Petróleo*, 4 (4).
- SARS, G.O. (1872). – Undersøgelser over Hardangerfjordens Fauna. – *Forhandlinger i Videnskasselskabet i Kristiania*, 1871, 246-255.
- SARS, M. (1869). – Fortsatte Bemaerkinger over det Dyriske Livs Udbredning i Havets Dybder. – *Forhandlinger i Videnskasselskabet i Kristiania*, 1868, 246-275.
- SCHAFFER, C.T., COLE, F.E. & CARTER, L. (1983). – Paraecology of bathyal zone arenaceous foraminifera genera and species associations off northeast Newfoundland. – *Proceedings of the 1st Workshop Arenaceous Foraminifera*. Continental Shelf Institute, Trondheim, Publ. 108, 133-147.
- SCHERER, R.P. & KOÇ, N. (1996). – Late Paleogene diatom biostratigraphy and paleoenvironments of the northern Norwegian Sea. – *Proceedings of the Ocean Drilling Program, Scientific Results*, 151, 75-99.
- SCHREIBER, O. (1980). – Bentonische Foraminiferen der Pemberger-Folge (Oberkreide) von Klein-Sankt Paul am Krappfeld (Karnten). – *Beiträge zur Paläontologie Österreich*, 7, 119-237. Wien.

- SCHRÖDER, C.J. (1986a). – Deep-water arenaceous foraminifera in the northwest Atlantic Ocean. – *Canadian Technical Report of Hydrography and Ocean Sciences*, 71, 191 pp.
- SCHRÖDER, C.J. (1986b). – Changes in benthic foraminifer assemblages across the Holocene/Pleistocene boundary, Sites 619, 620, 621, 622, and 624, Deep Sea Drilling Project Leg 96. – *Initial Reports of the Deep Sea Drilling Project*, 96, 631- 642.
- SCHRÖDER, C.J. (1988). – Subsurface preservation of arenaceous foraminifera in the northwest Atlantic Ocean. – *In: Proceedings to the Second Workshop on Agglutinated Foraminifera*, Vienna 1986. – *Abhandlungen der Geologischen Bundesanstalt*, 41, 325-336.
- SCHRÖDER-ADAMS, C.J. (1990). – High latitude agglutinated foraminifera: Prydz Bay (Antarctica) vs. Lancaster Sound (Canadian Arctic). – *In: Hemleben, C., Kaminski, M.A., Kuhnt, W., and Scott, D.B. (eds), Paleoecology, Biostratigraphy, Paleooceanography and Taxonomy of agglutinated foraminifera*. NATO ASI C, 327, 315-343. Kluwer Academic Press.
- SCHRÖDER-ADAMS, C.J. & MCNEIL, D.H. (1994a). – New paleoenvironmentally important species of agglutinated foraminifera from the Oligocene and Miocene of the Beaufort Sea, Arctic Canada. – *Journal of Foraminiferal Research*, 24, 178-190.
- SCHRÖDER-ADAMS, C.J. & MCNEIL, D.H. (1994b). – Oligocene to Miocene agglutinated foraminifera in deltaic and deep-water facies of the Beaufort-MacKenzie Basin. – *Geological Survey of Canada Bulletin*, 477, 67 pp.
- SCHUBERT, R.J. (1902). – Neue und interessante Foraminiferen aus dem südtiroler Altertüm. – *Beiträge zur Paläontologie und Geologie Österreich-Ungarns und des Orients*, 14, 9-26. Wien.
- SCHULTZE, F.E. (1875). – *Zoologische Ergebnisse der Nordseefahrt vom 21 Juli bis 9 September 1872; 1. Rhizopoden*. – *Comm. Wiss. Untersuchung Deutsch. Meere Kiel, Jahresber.*, Berlin, Deutschland, 1875, Jahrg. 2-3.
- SCOTT, D., GRADSTEIN, F.M., SCHAFFER, C.T., MILLER, A. & WILLIAMSON, M.A. (1983). – The Recent as a key to the Past: Does it apply to agglutinated foraminiferal assemblages. – *Proceedings of the 1st Workshop Arenaceous Foraminifera*. Continental Shelf Institute, Trondheim, Publ. 108, pp. 147-157.
- SCOTT, G.H. (1961). – Contribution to the knowledge of *Rzehakina* Cushman (Foraminifera) in New Zealand. – *New Zealand Journal of Geology and Geophysics*, 4, 3-43.
- SEIBOLD, E. & SEIBOLD, J. (1955). – Revision der Foraminiferen Bearbeitung C.W. Gümbel (1862) aus den Streitberg Schwamm Mergeln (Oberfranken, Unterer Malm). – *Neues Jahrbuch für Paläontologie, Abhandlungen*, 101, 91-134.
- SEIGLIE, G.A. & BAKER, M.B. (1987). – *Ammonia rosea*, new genus and new species of deep-water Trochamminidae, Foraminifera. – *Proceedings of the Biological Society of Washington*, 100 (3), 490-493.
- SEIGLIE, G.A. & BAKER, M.B. (1987). – Duquepsamminiidae, a new family, and *Duquepsammia*, a new genus of agglutinated foraminifera. – *Micropaleontology*, 33, 263-266.
- SEROVA, M.Ya. (1969). – Comparison of characteristics of Rzehakinidae in the Carpathian region and Pacific Province. – *Rocznik Polskiego Towarzystwa Geologicznego*, 39, 225-240.
- SEROVA, M.Ya. (1987). – *Foraminifery i biostratigrafia Severnoy Patsifiki na rubiezhie Mela i Paleogena*. – Izdatelstvo "Nauka", Moscow. 144 pp. + 16 pls.
- SEROVA, M.YA., DANILESKO, L.A., KOLYADA, A.A. & PETRINA, N.M., 1980. – Raschlenenie vyerkhnelovykh i paleogenovykh otlozhenii Koryakskogo Nagor'ya. – *Izvestiya Akademii Nauk SSSR, Seriya Geologicheskaya*, 1980 (10), 73-83.
- Shipboard Scientific Party (1987). – Site 643. – *Proceedings of the Ocean Drilling Program, Initial Reports*, 454-615.
- SIDEBOTTOM, H. (1918). – Report on the Recent Foraminifera dredged off the east coast of Australia, HMS "DART" Station 19 (14 May, 1895), Lat. 29°22'S., Long. 153°51'E., 465 fathoms, Pteropod ooze. – *Journal of the Royal Microscopical Society*, 1918, 1-25.
- SKINNER, H.C. (1961) – Revision of "*Protonina difflugiformis*". – *Journal of Paleontology*, 35, 1238-1240.
- ŚLĄCZKA, A. (1993). – The Geology of the External Carpathians: The classic localities revisited. – *In: Kaminski, M.A., Geroch, S. & Kaminski, D. (eds), The Origins of Applied Micropaleontology: The School of Jozef Grzybowski*. Grzybowski Foundation Special Publication, 1, 225-237. Alden Press, Oxford.
- ŚLĄCZKA, A. & KAMINSKI, M.A. (1998). – *Guidebook to Excursions in the Polish Flysch Carpathians - Field excursions for Environmental Geoscientists*. – Grzybowski Foundation Special Publication, 6, 173 pp.
- ŚLĄCZKA, A., GEROCH, S. & KOSZARSKI, L. (1993). – *IV International Workshop on Agglutinated Foraminifera Excursion Guidebook, Polish Flysch Carpathians*. – Grzybowski Foundation Special Publication, 2, 54 pp.

- SLITER, W.V. (1968). – Upper Cretaceous foraminifera from southern California and northwestern Baja California, Mexico. – *University of Kansas Paleontological Contributions*, ser. 49, 7, 1-141. Lawrence, Kansas.
- SLITER, W.V. (1971). – Predation on benthic foraminifera. – *Journal of Foraminiferal Research*, 1, 20-28.
- SLITER, W.V. & BAKER, R.A. (1972). – Cretaceous bathymetric distribution of benthic foraminifers. – *Journal of Foraminiferal Research*, 2, 167-183.
- SOLIMAN, H.A. (1972). – New Upper Cretaceous foraminifera from Soviet Carpathian (USSR). – *Revue de Micropaleontologie*, 15, 35-44.
- SOKOLOVA, M.N. (1997). – Trophic structure of abyssal macrobenthos. – In: Gebruk, A.V., Southward, A.J. & Tyler, P.A. (eds), *Advances in Marine Biology*, 32, 427-525. Academic Press, London.
- SOKOLOVA, M.N. VINOGRADOVA, N.G. & BURMISTROVA, I.I. (1995). – "Rain of dead bodies as a trophic source for the ultraabyssal macrobenthos in the Orkney Trench. – *Okeanologiya*, 35 (4), 587-591 [in Russian].
- SPEIJER, R. (1994). – Extinction and recovery patterns in benthic foraminiferal paleocommunities across the Cretaceous/Paleogene and Paleocene/Eocene boundaries. – *Geologica Ultraiectina*, 124, 191 pp.
- STAESCHE, K. & HILTERMANN, H. (1940). – Mikrofaunen aus dem Tertiär Nordwestdeutschlands. – *Abhandlungen der Reichsstelle für Bodenforschung, neue Folge*, 201, 26 pp. + 52 pls.
- STAINFORTH, R.M. (1952). – Ecology of arenaceous foraminifera. – *The Micropaleontologist*, 6, 42-44.
- STRÁNÍK, Z. & HANZLÍKOVÁ, E. (1963). – Geologická studie smilenského okna. – *Sborník Ústředního Ústavu Geologického, oddíl geologický*, 28, 457 - 495.
- STRUCK, U. (1995). – Stepwise postglacial migration of benthic foraminifera into the abyssal northeastern Norwegian Sea. – *Marine Micropaleontology*, 26, 201-213.
- SUBBOTINA, N.N. (1950). – Mikrofauna I stratigrafiya El'burganskogo Gorizonta I Gorizonta Goryachego Klyucha [Microfauna and stratigraphy of the El'burganskii horizon and the Goryachii Klyuch horizon]. – *Mikrofauna SSSR, vol 4. Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Instituta VNIGRI*, 51, 5-112+ 5 pls.
- SUBBOTINA, N.N. (1960). – Mikrofauna oligotsenovykh I Miotsenovykh otlozhenii r. Vorotyshche (Predkarpat'e [Microfauna of the Oligocene and Miocene deposits of the Vorotyshche River (Fore-Carpathians)]. – *Mikrofauna SSSR, vol. 11. – Trudy Vsesoyuznogo Nauchno- Issledovatel'skogo Geologorazvedochnogo Instituta VNIGRI*, 153, 157-243 + 10 pls.
- SULEYMANOV, I.S. (1959). – O novom rodye i vidye foraminifer iz sem. Ammodiscidae [On a new genus and species of foraminifera from the family Ammodiscidae]. – *Doklady Akademi Nauk UzSSR*, 1959 (7), 19-20.
- SULEYMANOV, I.S. (1960). – Noviy podrod i dva novykh vida iz semeystva Ammodiscidae [A new subgenus and two new species from the family Ammodiscidae]. – *Doklady Akademi Nauk UzSSR*, 1960 (2), 18-20.
- SULEYMANOV, I.S. (1963). – Noviy rod i dva novykh vida iz semeystva Ammodiscidae [A new genus and two new species from the family Ammodiscidae]. – *Doklady Akademi Nauk UzSSR*, 1963 (11), 41-43.
- SVAVARSSON, J., GUDMUNDSSON, G. & BRATTEGARD, T. (1993). – Feeding by *Asellote isopods* (Crustacea) on Foraminifers (Protozoa) in the deep sea. – *Deep-Sea Research Part I-Oceanographic Research Papers*, 40 (6), 1225-1239
- SZCZUCHURA, J. & POŻARYSKA, K. (1974). – Foraminiferida from the Paleocene of Polish Carpathians (Babice Clays). – *Paleontologica Polonica*, 31, 1-142.
- SZTRÁKOS, K. (1979). – La Stratigraphie, paléoécologie, paléogéographie et les foraminifères de l'Oligocène du nord-est de la Hongrie. – *Cahiers de Micropaléontologie*, 3, 7-95 + 35 pls.
- SZTRÁKOS, K. (1982). – Les foraminifères de la Marne de Buda et la limite Éocène-Oligocène en Hongrie. – *Cahiers de Micropaléontologie*, 4, 3-48 +39 pls.
- TAKAYANAGI, Y. (1960). – Cretaceous foraminifera from Hokkaido, Japan. – *Science Reports of the Tohoku University, Sendai, Japan, second series (Geology)*, 32 (1), 154 pp + 11 pls.
- TAPPAN, H. (1960). – Cretaceous biostratigraphy of northern Alaska. – *American Association of Petroleum Geologists Bulletin*, 44, 273-297.
- TEN DAM, A. & SIGAL, J. (1950). – Some new species of foraminifera from the Dano-Montian of Algeria. – *Contributions from the Cushman Foundation for Foraminiferal Research*, 1, 31-37.
- THALMANN, H.E. (1932). – *Nonion Jarvisi* nom. nov. and *Trochammina Kellettae* nom. nov. – *Ecologiae geologicae Helvetiae*, 25, 312-313.
- THALMANN, H.E. (1947). – Index to new genera, species and varieties of foraminifera for the year 1945 with supplements for the period 1939-1944, and addenda (1942-1945). – *Journal of Paleontology*, 21, 355-395.

- THALMANN, H.E. (1949). – Regional-stratigraphische Verbreitung der Foraminiferen-Gattung *Rzehakina* Cushman, 1927. – *Eclogae Geologicae Helvetiae*, 42 (2), 506 - 507.
- THEYER, F. (1971). – Size-depth variation in *Cyclammina cancellata* Brady, Peru-Chile Trench area. – In: Ried, J.L. (Ed.), *Biology of the Antarctic Seas*. Antarctic Research Series, 15, 309-313. American Geophysical Union.
- THIES, A. (1991). – Die Benthos-Foraminiferen im Europäischen Nordmeer. – *Berichte aus dem Sonderforschungsbereich 313*, 31, 97 pp. + 20 pl.
- THOMAS, E. (1985). – Late Eocene to Recent deep-sea benthic foraminifers from the Central Equatorial Pacific. – *Initial Reports of the Deep Sea Drilling Project*, 85, 655-694.
- THOMAS, E. & SHACKLETON, N.J. (1996). – The Paleocene–Eocene benthic foraminiferal extinction and stable isotope anomalies. – In: Knox, R.W., Corfield, R.M. & Dunar, R.E. (eds), *Correlation of the Early Paleogene in Northwest Europe*. Geological Society Special Publication, 101, 401-441.
- THOMAS, F.C. (1985). – *Lower Scotian Slope benthic foraminiferal faunas past and present, with taxonomic outline* [M.Sc. Thesis]. – Dalhousie University, Halifax. 159pp.
- TJALSMA, R.C. (1976) – Cenozoic foraminifera from the South Atlantic, DSDP Leg 36. – *Initial Reports of the Deep Sea Drilling Project*, 36, 493-517.
- TJALSMA, R.C. & LOHMANN, G.P. (1983). – Paleocene-Eocene bathyal and abyssal benthic foraminifera from the Atlantic Ocean. – *Micropaleontology Special Publication*, 4, 1-90.
- TODD, R. (1966). – Smaller foraminifera from Guam. – *US Geological Survey Professional Paper*, 403-I, 141 pp.
- TODD, R. & KNICKER, H.T. (1952). – An Eocene foraminiferal fauna from the Agua Fresca Shale of the Magallanes Province, southernmost Chile, – *Cushman Foundation for Foraminiferal Research Special Publication*, 1, 28 pp.
- TRUJILLO, E.F. (1960). – Upper Cretaceous foraminifera from near Redding, Shasta County, California. – *Journal of Paleontology*, 34, 290-346.
- TUCHOLKE, B.E. & VOGT, P.R. (1979). – Western North Atlantic: sedimentary evolution and aspects of tectonic history. – *Initial Reports of the Deep Sea Drilling Project*, 43, 791-825.
- TURENKO, T.V. (1983). – New foraminiferal species of the Family Rzehakinidae from the Upper Cretaceous of Sakhalin. – *Paleontologicheskii Zhurnal*, 1983(3), 13-21.
- UCHIO, T. (1960). – Ecology of living benthonic foraminifera from the San Diego, California area. – *Cushman Foundation for Foraminiferal Research Special Publication*, 5, 72 pp.
- VAN COUVERING, J.A., AUBRY, M.P., BERGGREN, W.A., BUJAK, J.P., NAESER, C.W. & WIESER, T. (1981). – The terminal Eocene event and the Polish connection. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, 36, 321-362.
- VAN DEN AKKER, T.J.H.A., KAMINSKI, M.A., GRADSTEIN, F.M. & WOOD, J., (2000). – Campanian to Palaeocene biostratigraphy and palaeoenvironments in the Foula Sub-basin, west of the Shetland Islands, UK. – *Journal of Micropalaeontology*, 19 (1), 23-43.
- VAN DER ZWAAN, G.J., DUIJNSTEE, I.A.P., DEN DULK, M., ERNST, S.R., JANNINK, N.T. & KOUWENHOVEN, T.J. (1999). – Benthic foraminifers: proxies or problems? A review of paleocological concepts. – *Earth Science Reviews*, 46 (1-4), p. 213-236.
- VAŠIČEK, M. (1947). – Poznámky k mikrobiostratigrafii magurského flyše na Moravě. – *Věstník Státního Geologického Ústavu Československé Republiky*, 22, 235-256.
- VERDENIUS, J.G. & VAN HINTE, J.E. (1983). – Central Norwegian-Greenland Sea: Tertiary arenaceous foraminifera, biostratigraphy and environment. – Proceedings of the First Workshop Arenaceous Foraminifera, 7-9 Sept. 1981. *Continental Shelf Institute Publication*, 108, 173-224.
- VOLAT, J.L., HUGO, B. & BIGNOUMBA-ILOGUE, J. (1996). – Foraminifères arénacés du Crétacé supérieur du Gabon. – *Bulletin des Centres de Recherches Exploration-Production Elf Aquitaine*, 20, (1), 229-275.
- VOLOSHINOVA, N.A. & BUDASHEVA, A.I. (1961). – Lituolids and trochamminids from the Tertiary deposits of Sakhalin Island and the Kamchatka Peninsula (in Russian). – In: *Microfauna of the USSR, Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Instituta VNIGRI*, 170, 170-272. Leningrad.
- VOLOSHINOVA, N.A., KUZNETSOVA, V.N. & LEONENKO, L.S. (1970). – Foraminifery Neogenovykh otlozhenii Sakhalina. – *Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Geologorazvedochnogo Instituta VNIGRI*, 284, 304 pp.
- VIALOV, O.S. (1966). – Zamechaniya o foraminiferakh s kremnevoy rakovinoy [Remarks on foraminifera with siliceous test]. – *Paleontologicheskii Sbornik, L'vov*, 3, 3-11. Lvov.

- VIALOV, O.S. (1966). – O krupnykh kremnistykh foraminiferakh Silicifera iz Verkhnemelovogo flisha Karpat [Large siliceous foraminifer Silicifera from the Upper Cretaceous flysch of the Carpathians]. – *Paleontologicheskii Sbornik, L'vov*, 3, 27-36
- VIALOV, O.S. & DABAGIAN, N.V. (1967). – Notes on some Paleogene Foraminifera from Grzybowski's collection (English summary). – *Paleontologicheskii Sbornik, L'vov*, 4, 24-34.
- VIALOV, O.S., GAVURA, S.P., DANYSH, V.V. *et al.* (1988). – *Stratotypy melovykh i paleogenovykh otlozhenii Ukrainiskikh Karpat. [Statotypes of Cretaceous and Paleogene deposits of the Ukrainian Carpathians]*. – Naukova Dumka (Kiev), 204 pp.
- VIALOV, O.S., MASLUN, N.V. & SOVCHIK, Ya.V. (1984). – K korrelyatsii paleogenogo flisha ukrainskikh i rumynskikh karpat [On the correlation of the flysch of the Ukrainian and Romanian Carpathians]. – *Anuarul institutului de Geologie si Geofizica Bucharest*, 59, 277-285.
- WEBB, P.N. (1972). – Paleocene foraminifera from Wangaloa and Dunedin. – *New Zealand Journal of Geology and Geophysics*, 16, 109-157.
- WEBB, P.N. (1975). – Paleocene foraminifera from DSDP Site 283, South Tasman Basin. – *Initial Reports of the Deep Sea Drilling Project*, 29, 833-843.
- WEIDICH, K.F. (1987). – Das Ultrahelvetikum von Liebenstein (Allgau) und seine Foraminiferenfauna. – *Zitteliana*, 15, 193-217.
- WEIDICH, K.F. (1990). – Die kalkalpine Unterkreide und ihre Foraminiferenfauna. – *Zitteliana*, 17, 3-312.
- WENGER, W.F. (1987). – Die Foraminiferen des Miozäns der bayerischen Molasse und ihre stratigraphische sowie paläogeographische Auswertung. – *Zitteliana*, 16, 173-340.
- WHITE, M.P. (1928a). – Some index foraminifera of the Tampico Embayment area of Mexico. – *Journal of Paleontology*, 2, 177-215.
- WHITE, M.P. (1928b). – Some index foraminifera of the Tampico Embayment area of Mexico (Part 2). – *Journal of Paleontology*, 2, 280-317.
- WHITE, M.P. (1929). – Some index foraminifera of the Tampico Embayment area of Mexico (Part 3). – *Journal of Paleontology*, 3, 30-58.
- WHITMARSH, R.B., BESLIER, M.O., WALLACE, P.J., *et al.* (1998). – *Proceedings of the Ocean Drilling Program, Initial Reports*, 173, 493 pp.
- WICKENDEN, R.T.D. (1932). – New species of foraminifera from the Upper Cretaceous of the Prairie Provinces. – *Trans. Royal Soc. Canada ser. 3*, 26, 85-91.
- WIDMARK, J.G.V. (1997). – Deep-sea benthic foraminifera from Cretaceous-Paleogene boundary strata in the South Atlantic - taxonomy and paleoecology. – *Fossils & Strata*, 43, 94 pp.
- WIESNER, H. (1931). – Die Foraminiferen der Deutsche Südpolar Expedition 1901-1903. – *Deutsche Südpolar Expedition*, vol 20, Zoologie, 165 pp + 24 pls. Walter de Gruyter & Co., Berlin.
- WIGHTMAN, W.G. & KUHN, W. (1992). – Biostratigraphy and paleoecology of Late Cretaceous abyssal agglutinated foraminifers from the western Pacific Ocean (Deep Sea Drilling Project Holes 196A and 198A and Ocean Drilling Program Holes 800A and 801A). – *Proceedings of the Ocean Drilling Program, Scientific Results*, 129, 247-264
- WILLIAMSON, M.A. 1983. – *Benthic foraminiferal assemblages on the continental margin off Nova Scotia, a multivariate approach*. – Ph.D. Thesis, Dalhousie University.
- WINKLER, W. (1984). – *Rhabdammina* fauna: What relation to turbidites? Evidence from the Gurnigel-Schlieren Flysch. – *In: Oertli, H.J. (ed.), Benthos '83, 2nd Int. Symp. Benthic Foraminifera*, Pau (France), April 11-15, 1983. Elf Aquitaine, ESSO REP and TOTAL CFP, Pau & Bourdeaux, pp. 611-617.
- WOLLENBURG, J.E. & KUHN, W. (2000). – The response of benthic foraminifers to carbon flux and primary production in the Arctic Ocean. – *Marine Micropaleontology*, 40, 189-231.
- WOOD, A. (1949). – The structure of the wall of the test in foraminifera: its value in classification. – *Quarterly Journal of the Geological Society of London*, 104, 229-255.
- YOUNG, F.G. & McNEIL, D.H. (1984). – Cenozoic stratigraphy of the MacKenzie Delta, Northwest Territories. – *Geological Survey Canada Bulletin*, 336, 63 pp.
- ZHENG, S. & FU, Z. (2001). – *Fauna Sinica, Phylum Granuloreticulosa, Class Foraminifera, Agglutinated Foraminifera*. Science Press, Beijing, 788 pp. + 122 pls.

